

BRAID MONODROMY COMPUTATION OF REAL SINGULAR CURVES

KAPLAN SHMUEL, LIBERMAN ERAN, AND TEICHER MINA

ABSTRACT. We generalize the Moishezon Teicher algorithm that was suggested for the computation of the braid monodromy of an almost real curve. The new algorithm suits a larger family of curves, and enables the computation of braid monodromy not only of caspidal curves, but of general algebraic curves, with some non simple singularities. Moreover, it works also when in the fiber the curve admits any number of imaginary points. We also provide two examples of how to use the generalized algorithm.

INTRODUCTION

Classification of surfaces is a major area of study in algebraic geometry. Hence, different methods for achieving this goal have grown to be very important. It was discovered that branch curves of hypersurfaces are invariant in some sense for those hypersurfaces, and the braid monodromy was suggested as an invariant for the branch curves.

The braid monodromy is an important tool in order to compute invariants of curves and surfaces, for example it is used as one way to compute the fundamental group of the complement of a curve. This invariant was used by Chisini [5], Kulikov [10] and Kulikov-Teicher [11] in order to distinguish between connected components of the moduli space of surfaces of general type.

Many previous works were done using the braid monodromy in order to classify surfaces, branch curves and in order to compute the fundamental groups of the complement of curves (for example see [6, 1, 2] among many others). However, there is still a growing need of computed examples. Until now, most examples were computed on caspidal curves with only simple singularities and limited number of imaginary points.

Date: February 1, 2008.

This paper is part of the First and Second authors PhD. Thesis in Bar-Ilan university.

Partially supported by the Emmy Noether Research Institute for Mathematics, Bar-Ilan University and the Minerva Foundation, Germany, and by the Excellency Center “Group theoretic methods in the study of algebraic varieties” of the National Science Foundation of Israel, and by Eager, European Algebraic Geometry Research.

In this paper we improve the algorithm given in [9] for the computation of braid monodromy of an almost real curve. The improved algorithm works on general algebraic curves, it enables to consider other singularities which were not considered in the past, where some of the singularities are not simple, and it also enables the computation with any number of complex points in any fiber above the real line.

The paper is organized as follows: In Section 1 we give preliminaries and definitions. Section 2 is devoted to the computations of the local braid monodromy of the new types of singular points. In Section 3 we present a new model for the fiber of the curve above a given point. We introduce the algorithm, which calculates the braid monodromy, and prove it's steps in Section 4. Finally, in Section 5 we give two examples for computations according to the algorithm.

1. PRELIMINARIES

We assume that the reader is familiar with the braid group and the braid monodromy subjects (see for example [3, 4, 7, 9]); Therefore, we give here only notations and some of the basic definitions involved.

Let S be a curve in $\mathbb{C}^2$. Denote by $\pi_1 : S \rightarrow \mathbb{C}$ and by $\pi_2 : S \rightarrow \mathbb{C}$ the projection to the first and second coordinate respectively, defined in the obvious way. Let n to be the degree of S with respect to π_1 . For $x \in \mathbb{C}$ we denote $K(x)$ the second coordinate of the points in S which are in the fiber above x (i.e., $K(x) = \pi_2(\pi_1^{-1}(x))$).

Let $N \subset \mathbb{C}$ be the set $N = N(S) = \{x \in \mathbb{C} \mid |K(x)| < n\} = \{x_1, \dots, x_p\}$. Take E to be a closed disc in $\mathbb{C}$ for which $N \subset E \setminus \partial E$. In addition take D to be a closed disc in $\mathbb{C}$ for which D contains all the points $\{K(x) \mid x \in E\}$. That means that when restricted to E , we have $S \subset E \times D$.

With these definitions we define the *braid monodromy*

Definition 1.1. *Let $M \in \partial E$ be the base point of $\pi_1(E \setminus N)$, and let σ be an element of $\pi_1(E \setminus N)$. To σ there are n lifts in S , each begins and ends at the points of $M \times K(M)$. Projecting these lifts using $\pi_2 : S \rightarrow \mathbb{C}$ we get n paths in D which begin and end at the points of $K(M)$. These paths induce a diffeomorphism of $\pi_1(D \setminus K(M))$ which is an element of the $\mathcal{MCG}$ of $\pi_1(D \setminus K(M))$ known to be the braid group B_n [4]. We call the homomorphism $\varphi : \pi_1(E \setminus N) \rightarrow B_n$ the braid monodromy of S with respect to $E \times D$, π_1 , and M .*

1.1. Real singular curves. We call the curve S a *real singular curve* if it is defined by a polynomial with real coefficients, and if the set $N(S) \subset \mathbb{R}$. We restrict ourselves to work with subset of real singular curves as follows:

Definition 1.2. Let $B = B(S) = \{x \in \mathbb{R} \mid \exists z_1, z_2 \in K(x) \setminus \mathbb{R} \text{ s.t. } \Im(z_1) = \Im(z_2)\}$.

Note that if $z \in K(x) \setminus \mathbb{R}$ so does its complex conjugate $\bar{z}$ (because, S is defined over $\mathbb{R}$) so when $x \in B$ we have at least two pairs of points z_1, z_2 and $\bar{z}_1, \bar{z}_2$ such that $\Im(z_1) = \Im(z_2)$, $\Im(\bar{z}_1) = \Im(\bar{z}_2)$ and $\Im(z_1) = -\Im(\bar{z}_1)$.

We work with curve S which satisfies the following conditions:

- (1) S is real singular.
- (2) For all $x \in N$ there is only one singular points of S with x as its first coordinate.
- (3) The set B is finite.
- (4) For all $x \in B$ the number of pairs $z_1, z_2 \in K(x) \setminus \mathbb{R}$ s.t. $\Im(z_1) = \Im(z_2)$ is 2.
- (5) $B \cap N = \emptyset$.

Our algorithm will apply to curves with the following types of singularities (looking at these singularities was motivated by the fact that we are interested mostly, but not only, in computing braid monodromy for branch curves of hypersurfaces).

- (1) a_1 : a branch point, topologically equivalent to $y^2 - x = 0$.
- (2) a_2 : a branch point, topologically equivalent to $y^2 + x = 0$.
- (3) b : a tangent point, topologically equivalent to $y(y - x^2) = 0$.
- (4) c : intersection of m non singular branches of S , transversal to each other.
- (5) d_1 : a cusp point, topologically equivalent to $y^2 - x^\nu = 0$, ($3 \leq \nu$).
- (6) d_2 : a cusp point, topologically equivalent to $y^2 + x^\nu = 0$, ($3 \leq \nu$).
- (7) d_3 : a cusp point, topologically equivalent to $y^3 - x^2 = 0$.
- (8) e_1 : two cusps intersecting at one point, topologically equivalent to $(y^2 + x^3)(y^2 - x^3) = 0$.
- (9) e_2 : two cusps intersecting at one point, topologically equivalent to $(y^3 + ax^2)(y^3 - bx^2) = 0$ ($a \neq b$), ($0 < a, b$).
- (10) f_1 : a cusp point intersecting a line, topologically equivalent to $y(y^2 - x^\nu) = 0$.
- (11) f_2 : a cusp point intersecting a line, topologically equivalent to $y(y^2 + x^\nu) = 0$.
- (12) g_1 : a cusp point intersecting a line, topologically equivalent to $(y - ax)(y^2 - x^\nu) = 0$.
- (13) g_2 : a cusp point intersecting a line, topologically equivalent to $(y - ax)(y^2 + x^\nu) = 0$.
- (14) g_3 : a cusp point intersecting a line, topologically equivalent to $(y)(y^3 - x^2) = 0$.

Remark 1.3.

- (1) *In our case topologically equivalent means that there is a deformation which transforms the given singular point in the curve S into singularity types defined above, without changing the multiplicity of any point with respect to the projection π_1 .*
- (2) *Without loss of generality we assume that $a > 0$ in the cases of singularity of type g_1 and g_2 since when $a < 0$ all constructions and proofs are similar.*
- (3) *The points of type d_1 and d_2 generalize the results previously obtained in [9] for the points a_1 and a_2 . We keep the notations of the points a_1 and a_2 in order to comply with previous works.*

The computation of the braid monodromy of the curve is done using computations of local braid monodromy at each singular point and combining the results. So our aim now is to describe the local braid monodromy at each type of singularity defined above.

2. COMPUTATION OF LOCAL BRAID MONODROMY

In this section we compute the local braid monodromy induced by each of the singularities defined above. We use the connection between the geometrical braid and its description using Artin generators [3, 4], by looking at the projection of the geometrical 3-dimensional braid to the xz plane keeping in mind which strings come above the other at intersection points of the projection. When two strings start or end at the same position we always rotate the braid clockwise by a very small angle in order to get a general position of the beginning and ending points of the strings.

Local braid monodromy of singular points of types a_1, a_2, b, c, d_1 were discussed in [8], [9] we will not repeat these computations here.

2.1. Local braid monodromy of singularity type d_2 . The computation here is similar to the one presented in [8][Proposition-Example VI.1.1]

Proposition 2.1. *Let S be the curve $y^2 + x^\nu$. Then, the local braid monodromy at the origin is generated by the braid σ_1^ν .*

Proof. Take E to be the unit disk, D be a closed disk sufficiently large such that $S|_E \subset E \times D$. $N = (0 + 0i) \in E$. We take $\Gamma = \{x = e^{2\pi it} \mid t \in [0, 1]\}$ to be the element of $\pi_1(E \setminus N, 1)$. Thus, when lifting Γ into S we get two paths:

$$\begin{aligned}\delta_1(t) &= (e^{2\pi it}, e^{\frac{\nu(2\pi it + \pi i)}{2}}) \\ \delta_2(t) &= (e^{2\pi it}, -e^{\frac{\nu(2\pi it + \pi i)}{2}})\end{aligned}$$

Where, $t \in [0, 1]$

Projecting them to D we get two paths:

$$a_1(t) = e^{\frac{\nu(2\pi it + \pi i)}{2}}$$

$$a_2(t) = -e^{\frac{\nu(2\pi it + \pi i)}{2}}$$

Which induce the braid σ_1' which transpose the position of two strings ν times, as described in Figure 1.




FIGURE 1. The curve of type d_2 where $\nu = 5$.

□

2.2. Local braid monodromy of singularity type d_3 .

Proposition 2.2. *Let S be the curve $y^3 - x^2$. Then, the local braid monodromy at the origin is generated by the braid $(\sigma_1\sigma_2)^2$.*

Proof. Take E to be the unit disk, D be a closed disk sufficiently large such that $S|_E \subset E \times D$. $N = (0 + 0i) \in E$. We take $\Gamma = \{x = e^{2\pi it} \mid t \in [0, 1]\}$ to be the element of $\pi_1(E \setminus N, 1)$. Thus, when lifting Γ into S we get 3 paths:

$$\delta_k(t) = (e^{2\pi it}, e^{\frac{4}{3}\pi it + \frac{2}{3}\pi ik})$$


Where, $t \in [0, 1]$ and $k = 0, 1, 2$.

Projecting them to D we get 3 paths:

$$a_k(t) = e^{\frac{4}{3}\pi it + \frac{2}{3}\pi ik}$$

Where $k = 0, 1, 2$.

Which induce the braid $(\sigma_1\sigma_2)^2$, see Figure 2.

FIGURE 2. The curve of type d_3 .

□

2.3. Local braid monodromy of singularity type e_1 .

Proposition 2.3. *Let S be the curve $(y^2 + x^3)(y^2 - x^3)$. Then, the local braid monodromy at the origin is generated the braid $(\sigma_2\sigma_1\sigma_3)^6$.*

Proof. Take E to be the unit disk, D be a closed disk sufficiently large such that $S|_E \subset E \times D$. $N = (0 + 0i) \in E$. We take $\Gamma = \{x = e^{2\pi it} \mid t \in [0, 1]\}$ to be the element of $\pi_1(E \setminus N, 1)$. Thus, when lifting Γ into S we get 4 paths:

$$\delta_1(t) = (e^{2\pi it}, e^{3\pi it})$$

$$\delta_2(t) = (e^{2\pi it}, -e^{3\pi it})$$

$$\delta_3(t) = (e^{2\pi it}, e^{3\pi it + \frac{\pi}{2}i})$$

$$\delta_4(t) = (e^{2\pi it}, -e^{3\pi it + \frac{\pi}{2}i})$$

Where, $t \in [0, 1]$

Projecting them to D we get two paths:


$$a_1(t) = e^{3\pi it}$$

$$a_2(t) = -e^{3\pi it}$$

$$a_3(t) = e^{3\pi it + \frac{\pi}{2}i}$$

$$a_4(t) = -e^{3\pi it + \frac{\pi}{2}i}$$

Which induce the braid $(\sigma_2\sigma_1\sigma_3)^6$, as can be seen in Figure 3.

FIGURE 3. The curve of type e_1 .

□

2.4. Local braid monodromy of singularity type e_2 .

Proposition 2.4. *Let S be the curve $(y^3 + x^2)(y^3 - x^2)$. Then, the local braid monodromy at the origin is generated by the braid $(\sigma_2\sigma_4\sigma_1\sigma_3\sigma_5)^4$.*

Proof. Take E to be the unit disk, D be a closed disk sufficiently large such that $S|_E \subset E \times D$. $N = (0 + 0i) \in E$. We take $\Gamma = \{x = e^{2\pi it} \mid t \in [0, 1]\}$ to be the element of $\pi_1(E \setminus N, 1)$. Thus, when lifting Γ into S we get 6 paths:

$$\delta_{1,k}(t) = (e^{2\pi it}, e^{\frac{4\pi it}{3} + \frac{2\pi ik}{3}})$$

$$\delta_{2,k}(t) = (e^{2\pi it}, e^{\frac{4\pi it}{3} + \frac{\pi}{3} + \frac{2\pi ik}{3}})$$


Where, $t \in [0, 1]$, and $k = 1, \dots, 3$.

Projecting them to D we get two paths:

$$a_{1,k}(t) = e^{\frac{4\pi it}{3} + \frac{2\pi ik}{3}}$$

$$a_{2,k}(t) = e^{\frac{4\pi it}{3} + \frac{\pi}{3} + \frac{2\pi ik}{3}}$$

Which induce the braid $(\sigma_2\sigma_4\sigma_1\sigma_3\sigma_5)^4$, as can be seen in Figure 4.

FIGURE 4. The curve of type e_2 .

□

2.5. Local braid monodromy of singularity type f_1 .

Proposition 2.5. *Let S be the curve $(y^2 - x^\nu)y$. Then, the local braid monodromy at the origin is generated by the braid $(\sigma_2\sigma_1\sigma_2)^\nu$.*

Proof. Take E to be the unit disk, D be a closed disk sufficiently large such that $S|_E \subset E \times D$. $N = (0 + 0i) \in E$. We take $\Gamma = \{x = e^{2\pi it} \mid t \in [0, 1]\}$ to be the element of $\pi_1(E \setminus N, 1)$. Thus, when lifting Γ into S we get Three paths:

$$\delta_1(t) = (e^{2\pi it}, e^{\nu\pi it})$$

$$\delta_2(t) = (e^{2\pi it}, -e^{\nu\pi it})$$

$$\delta_3(t) = (e^{2\pi it}, 0)$$

Where, $t \in [0, 1]$.

Projecting them to D we get three paths:

$$a_1(t) = e^{\nu\pi it}$$

$$a_2(t) = -e^{\nu\pi it}$$

$$a_3(t) = 0$$

Which induce the braid $(\sigma_2\sigma_1\sigma_2)^\nu$, as can be seen in Figure 5.




FIGURE 5. The curve of type f_1 where $\nu = 5$.

□

2.6. Local braid monodromy of singularity type f_2 .

Proposition 2.6. *Let S be the curve $(y^2 + x^\nu)y$. Then, the local braid monodromy at the origin is generated by the braid $(\sigma_2\sigma_1\sigma_2)^\nu$.*

Proof. The proof is similar to the proof of Proposition 2.5.

□

2.7. Local braid monodromy of singularity type g_1 .

Proposition 2.7. *Let S be the curve $(y^2 - x^\nu)(y - ax)$. Then, the local braid monodromy at the origin is generated by the braid $(\sigma_2\sigma_1\sigma_2^\nu\sigma_1\sigma_2)$.*

Proof. Without loss of generality we may choose $a = 2$, this will help us to avoid multiple intersections between the line and the cusp inside the unit disk. Take E to be the unit disk, D be a closed disk sufficiently large such that $S|_E \subset E \times D$ (in case where $a \neq 2$ we may need to take E to be a smaller disk). $N = (0 + 0i) \in E$. We take $\Gamma = \{x = e^{2\pi it} \mid t \in [0, 1]\}$ to be the element of $\pi_1(E \setminus N, 1)$. Thus, when lifting Γ into S we get Three paths:

$$\begin{aligned}\delta_1(t) &= (e^{2\pi it}, e^{\nu\pi it}) \\ \delta_2(t) &= (e^{2\pi it}, -e^{\nu\pi it}) \\ \delta_3(t) &= (e^{2\pi it}, 2e^{2\pi it})\end{aligned}$$

Where, $t \in [0, 1]$.

Projecting them to D we get three paths:

$$\begin{aligned}a_1(t) &= e^{\nu\pi it} \\ a_2(t) &= -e^{\nu\pi it} \\ a_3(t) &= 2e^{2\pi it}\end{aligned}$$

Which induce the braid $(\sigma_2\sigma_1\sigma_2^\nu\sigma_1\sigma_2)$, as can be seen in Figure 6.




FIGURE 6. The curve of type g_1 where $\nu = 5$.

□

2.8. Local braid monodromy of singularity type g_2 .

Proposition 2.8. *Let S be the curve $(y^2 + x^\nu)(y - ax)$. Then, the local braid monodromy at the origin is generated by the braid $(\sigma_2\sigma_1\sigma_2^\nu\sigma_1\sigma_2)$.*

Proof. The proof is similar to the proof of Proposition 2.7.

□

2.9. Local braid monodromy of singularity type g_3 .

Proposition 2.9. *Let S be the curve $(y)(y^3 - x^2)$. Then, the local braid monodromy at the origin is generated by the braid $(\sigma_1\sigma_2\sigma_3\sigma_2)^2$.*

Proof. Take E to be the unit disk, D be a closed disk sufficiently large such that $S|_E \subset E \times D$. $N = (0 + 0i) \in E$. We take $\Gamma = \{x = e^{2\pi it} \mid t \in [0, 1]\}$ to be the element of $\pi_1(E \setminus N, 1)$. Thus, when lifting Γ into S we get 4 paths:

$$\delta_k(t) = (e^{2\pi it}, e^{\frac{4}{3}\pi it + \frac{2}{3}\pi ik})$$

$$\delta_4(t) = (e^{2\pi it}, 0)$$

Where, $t \in [0, 1]$ and Where $k = 0, 1, 2$.

Projecting them to D we get two paths:

$$a_k(t) = e^{\frac{4}{3}\pi it + \frac{2}{3}\pi ik}$$

$$a_4(t) = 0$$

Where $k = 0, 1, 2$. Which induce the braid $(\sigma_1\sigma_2\sigma_3\sigma_2)^2$, as can be seen in Figure 7.




FIGURE 7. The curve of type g_3 .

□

3. A MODEL FOR THE FIBER

3.1. Model definition. In the algorithm we use a model of the fiber above a real point x_0 . In this fiber we distinguish n (the degree of the projection to the first coordinate π_1) real and complex points.

Since the model should be diffeomorphic to D we take the model to be a disk $\tilde{D} \subset \mathbb{C}$ which is centered at the point $n+1$ and have radius $n+1$. The distinguished real points will be in the set $K' = \{1, \dots, n\}$, and the distinguished complex points will be in the set $K'' = \{n+1 \pm i, \dots, n+1 \pm \lceil \frac{n}{2} \rceil i\}$.

Usually, we shell assume that we have d (an even number) complex points and therefore, we identify in $\tilde{D}$ the set of points $K_d = \{1, \dots, (n-d), (n+1) \pm i, \dots, (n+1) \pm \frac{d}{2}i\}$.

For example, if $n = 9$ and $d = 4$ (i.e., we have 4 complex points), we get $K_4 = \{1, 2, 3, 4, 5, 10+i, 10+2i, 10-i, 10-2i\}$ and the model is $\tilde{D}$ with the set of points K_4 identified, as can be seen in figure 8




FIGURE 8. E_4

3.2. Half-twists. We need to define some braids using our model, hence we need to describe an element of the $\mathcal{MCG}$ of the disk $\tilde{D}$ punctured at the points K_d .

Let σ be a simple path in $\tilde{D}$ connecting two points of K_d which do not intersect any point of K_d besides its two end points. Take a small neighborhood U of σ and a small neighborhood U' of U , which contain from K_d only the two end points of σ . We define a diffeomorphism which switches the two end points of σ along the curve counterclockwise, and is the identity outside U' . We call the diffeomorphism constructed the *(positive) half-twist* defined by σ , and we denote it by $H(\sigma)$. Note, different diffeomorphisms defined on the same curve σ , using the above construction, will induce the same braid, hence half-twists are well defined.

Definition 3.1. Let σ be a non self intersecting path in $\tilde{D} \setminus \partial\tilde{D}$ starting and ending at two different points of K_d , which go through all the points of K_d . The points of K_d induce a partition of σ into $n-1$ consecutive paths $\sigma_1, \dots, \sigma_{n-1}$. We call the ordered system of half-twists $H(\sigma_1), \dots, H(\sigma_{n-1})$ a frame of B_n .

It is well known that if $(H(\sigma_1), \dots, H(\sigma_{n-1}))$ is a frame of B_n , then the braid group is generated by its elements, and the relations are just like the relations in Artin presentation for the braid group [8].

3.3. Diffeomorphisms in the model. We will need to define braids using the model $\tilde{D}$ of the punctured disk $D \setminus K(x)$. For this matter we will give here a

few diffeomorphisms which we will use as building blocks. Note that all diffeomorphisms are the identity on $\partial\tilde{D}$. Moreover, instead of giving tedious formal definition, we will describe a continuous family of diffeomorphisms (starting with the identity), by the action they induce on a closed or a finite set of points in the interior of $\tilde{D}$. The braid we define will be the last diffeomorphism in the family.

Definition 3.2. *Let $R^n \langle s, r \rangle$ be the diffeomorphism which rotates counterclockwise in an angle of $2\pi n$ a closed disk centered at s and of radius r .*

Definition 3.3. *$L \langle c, k \rangle$ is the diffeomorphism which maps the point $c \in \mathbb{C}$ to the point $k \in \mathbb{C}$ and the point $\bar{c}$ (the complex conjugate of c) to the point $\bar{k}$ (the complex conjugate of k), along a simple path (usually a straight line) which does not intersect K'' .*

Definition 3.4. *$M \langle (h, j), (k, l) \rangle$ is the diffeomorphism which maps all points in the real segment $[h, j]$ onto the real segment $[k, l]$ (note that $j - h = l - k$).*

Definition 3.5. *C is the diffeomorphism which maps the points along the straight line between $n + 1 + i$ to $n + 1 + \frac{d}{2}i$ onto the straight line between $n + 1 + 2i$ and $n + 1 + (\frac{d}{2} + 1)i$, and the straight line between $n + 1 - i$ to $n + 1 - \frac{d}{2}i$ onto the straight line between $n + 1 - 2i$ and $n + 1 - (\frac{d}{2} + 1)i$.*

Definition 3.6. *$A^n \langle c, r_1, r_2 \rangle$ is the diffeomorphism which rotates counterclockwise by angle of $2\pi n$ the annulus centered at c between the radiuses r_1 and r_2 .*

We now turn to define several diffeomorphisms, each one of them in connection with a specific type of point defined in Section 1. Note, that since usually Δ denotes a half-twist in the braid group the powers of the Δ will be twice the angles of rotations used in Definitions 3.2 and 3.6.

$$\Delta_{a_1}^{\frac{1}{2}} \langle k, l \rangle = L \langle n + 1 + i, k + \frac{1}{2} + \frac{i}{2} \rangle \circ M \langle (k, n - d), (k + 2, n - d + 2) \rangle \circ R^{\frac{1}{4}} \langle k + \frac{1}{2}, \frac{1}{2} \rangle \circ C^{-1}$$

$$\Delta_{a_2}^{\frac{1}{2}} \langle k, l \rangle = R^{\frac{1}{4}} \langle k + \frac{1}{2}, \frac{1}{2} \rangle \circ C \circ L \langle k + \frac{1}{2} + \frac{i}{2}, n + 1 + i \rangle \circ M \langle (k + 2, n - d), (k, n - d - 2) \rangle$$

$$\Delta_b^2 \langle k, l \rangle = R^1 \langle k + \frac{1}{2}, \frac{1}{2} \rangle$$

$$\Delta_c \langle k, l \rangle = R^{\frac{1}{2}} \langle \frac{k+l}{2}, \frac{l-k}{2} \rangle$$

$$\Delta_{d_1}^{\frac{\nu}{2}} \langle k \rangle = \begin{cases} L \langle n + 1 + i, k + \frac{1}{2} + \frac{i}{2} \rangle \circ M \langle (k, n - d), (k + 2, n - d + 2) \rangle \circ R^{\frac{\nu}{4}} \langle k + \frac{1}{2}, \frac{1}{2} \rangle \circ C^{-1} & \text{if } \nu \text{ is odd} \\ R^{\frac{\nu}{4}} \langle k + \frac{1}{2}, \frac{1}{2} \rangle & \text{if } \nu \text{ is even} \end{cases}$$

$$\Delta_{d_2}^{\frac{\nu}{2}} \langle k, l \rangle = \begin{cases} R^{\frac{\nu}{4}} \langle k + \frac{1}{2}, \frac{1}{2} \rangle \circ C \circ L \langle k + \frac{1}{2} + \frac{i}{2}, n + 1 + i \rangle \circ \\ M \langle (k + 2, n - d), (k, n - d - 2) \rangle & \text{if } \nu \text{ is odd} \\ \\ L \langle n + 1 + i, k + \frac{1}{2} + \frac{i}{2} \rangle \circ M \langle (k, n - d), (k + 2, n - d + 2) \rangle \circ \\ R^{\frac{\nu}{4}} \langle k + \frac{1}{2}, \frac{1}{2} \rangle \circ \\ M \langle (k + 2, n - d + 2), (k, n - d) \rangle \circ L \langle k + \frac{1}{2} + \frac{i}{2}, n + 1 + i \rangle & \text{if } \nu \text{ is even} \end{cases}$$

$$\Delta_{d_3}^{\frac{2}{3}} \langle k \rangle = L \langle n + 1 + i, k + \frac{1}{4} + \frac{\sqrt{3}}{4}i \rangle \circ M \langle (k, n - d), (k + 1, n - d + 1) \rangle \circ \\ R^{\frac{1}{3}} \langle k + \frac{1}{2}, \frac{1}{2} \rangle \circ M \langle (k + 1, n - d + 1), (k, n - d) \rangle \circ L \langle k + \frac{1}{4} + \frac{\sqrt{3}}{4}i, n + 1 + i \rangle$$

$$\Delta_{e_1}^{\frac{3}{2}} \langle k, l \rangle = L \langle n + 1 + i, k + \frac{1}{2} + \frac{i}{2} \rangle \circ R^{\frac{3}{4}} \langle k + \frac{1}{2}, \frac{1}{2} \rangle \circ \\ L \langle k + \frac{1}{2} + \frac{i}{2}, n + 1 + i \rangle$$

$$\Delta_{e_2}^{\frac{2}{3}} \langle k, l \rangle_{\langle a < b \rangle} = L \langle n + 1 + i, l - \frac{1}{4} + \frac{\sqrt{3}}{4}i \rangle \circ L \langle n + 1 + 2i, k + \frac{1}{4} + \frac{\sqrt{3}}{4}i \rangle \circ R^{\frac{1}{3}} \langle k + \frac{1}{2}, \frac{1}{2} \rangle \circ \\ L \langle k + \frac{1}{4} + \frac{\sqrt{3}}{4}i, n + 1 + 2i \rangle \circ L \langle l - \frac{1}{4} + \frac{\sqrt{3}}{4}i, n + 1 + i \rangle$$

$$\Delta_{e_2}^{\frac{2}{3}} \langle k, l \rangle_{\langle a > b \rangle} = L \langle n + 1 + i, k + \frac{1}{4} + \frac{\sqrt{3}}{4}i \rangle \circ L \langle n + 1 + 2i, l - \frac{1}{4} + \frac{\sqrt{3}}{4}i \rangle \circ R^{\frac{1}{3}} \langle k + \frac{1}{2}, \frac{1}{2} \rangle \circ \\ L \langle l - \frac{1}{4} + \frac{\sqrt{3}}{4}i, n + 1 + 2i \rangle \circ L \langle k + \frac{1}{4} + \frac{\sqrt{3}}{4}i, n + 1 + i \rangle$$

For the next diffeomorphism we need to define $\mathcal{D}$ as follows:

$$\mathcal{D} = L \langle n + 1 + i, k + 1 + i \rangle \circ M \langle (k + 1, n - d), (k + 3, n - d + 2) \rangle \circ \\ M \langle (k, k), (k + 1, k + 1) \rangle$$

$$\Delta_{f_1}^{\frac{\nu}{2}} \langle k, l \rangle = \begin{cases} \mathcal{D} \circ R^{\frac{\nu}{4}} \langle k + 1, 1 \rangle \circ C^{-1} & \text{if } \nu \text{ is odd} \\ \\ R^{\frac{\nu}{4}} \langle k + 1, 1 \rangle & \text{if } \nu \text{ is even} \end{cases}$$

$$\Delta_{f_2}^{\frac{\nu}{2}} \langle k, l \rangle = \begin{cases} R^{\frac{\nu}{4}} \langle k + 1, 1 \rangle \circ C \circ L \langle k + 1 + i, n + 1 + i \rangle \circ & \text{if } \nu \text{ is odd} \\ M \langle (k + 1, k + 1), (k, k) \rangle \circ \\ M \langle (k + 3, n - d), (k + 1, n - d - 2) \rangle \\ \\ \mathcal{D} \circ R^{\frac{\nu}{4}} \langle k + 1, 1 \rangle \circ \mathcal{D}^{-1} & \text{if } \nu \text{ is even} \end{cases}$$

$$\Delta_{g_1}^{\frac{\nu}{2}} \langle k, l \rangle = \begin{cases} M \langle (k + 1, n - d), (k + 3, n - d + 2) \rangle \circ & \text{if } \nu \text{ is odd} \\ L \langle n + 1 + i, k + 1 + \frac{i}{2} \rangle \circ \\ A^{\frac{1}{2}} \langle k + 1, \frac{3}{4}, \frac{5}{4} \rangle \circ R^{\frac{\nu}{4}} \langle k + 1, \frac{1}{2} \rangle \circ \\ M \langle (k - \frac{1}{2}, k + \frac{1}{2}), (k, k + 1) \rangle \circ C^{-1} \\ \\ M \langle (k, k), (k + \frac{1}{2}, k + \frac{1}{2}) \rangle \circ A^{\frac{1}{2}} \langle k + \frac{3}{2}, \frac{3}{4}, \frac{5}{4} \rangle \circ & \text{if } \nu \text{ is even} \\ R^{\frac{\nu}{4}} \langle k + \frac{3}{2}, \frac{1}{2} \rangle \circ M \langle (k + 1, k + 2), (k, k + 1) \rangle \circ \\ M \langle (k + \frac{5}{2}, k + \frac{5}{2}), (k + 2, k + 2) \rangle \end{cases}$$

$$\Delta_{g_2}^{\frac{\nu}{2}} \langle k, l \rangle = \begin{cases} M \langle (k, k), (k + \frac{1}{2}, k + \frac{1}{2}) \rangle \circ A^{\frac{1}{2}} \langle k + \frac{3}{2}, \frac{3}{4}, \frac{5}{4} \rangle \circ & \text{if } \nu \text{ is odd} \\ R^{\frac{\nu}{4}} \langle k + \frac{3}{2}, \frac{1}{2} \rangle \circ C \circ L \langle k + \frac{3}{2} + \frac{i}{2}, n + 1 + i \rangle \circ \\ M \langle (k + \frac{5}{2}, k + \frac{5}{2}), (k, k) \rangle \circ \\ M \langle (k + 3, n - d), (k + 1, n - d - 2) \rangle \\ \\ M \langle (k + 1, n - d), (k + 3, n - d + 2) \rangle \circ & \text{if } \nu \text{ is even} \\ L \langle n + 1 + i, k + 1 + \frac{i}{2} \rangle \circ \\ A^{\frac{1}{2}} \langle k + 1, \frac{3}{4}, \frac{5}{4} \rangle \circ R^{\frac{\nu}{4}} \langle k + 1, \frac{1}{2} \rangle \circ \\ L \langle k + 1 + \frac{i}{2}, n + 1 + i \rangle \circ \\ M \langle (k + 2, n - d + 2), (k, n - d) \rangle \end{cases}$$

$$\Delta_{g_3}^{\frac{2}{3}} \langle k, l \rangle = M \langle (k + 1, k + 1), (k + \frac{1}{2}, k + \frac{1}{2}) \rangle \circ L \langle n + 1 + i, k - \frac{1}{4} + \frac{\sqrt{3}}{4}i \rangle \circ \\ R^{\frac{1}{3}} \langle k, \frac{1}{2} \rangle \circ L \langle k - \frac{1}{4} + \frac{\sqrt{3}}{4}i, n + 1 + i \rangle \circ \\ M \langle (k + \frac{1}{2}, k + \frac{1}{2}), (k + 1, k + 1) \rangle$$

$$\Delta_{com}^1 \langle k, l \rangle = R^{\frac{1}{2}} \langle n + 1 + (k + \frac{1}{2})i, \frac{1}{2} \rangle \circ R^{-\frac{1}{2}} \langle n + 1 - (k + \frac{1}{2})i, \frac{1}{2} \rangle$$

4. DESCRIPTION OF THE ALGORITHM

In this section we develop the theory needed for the algorithm. We prove the theorems which form the heart of the algorithm, and at the end of this section we give the algorithm's steps.

4.1. definitions.

Definition 4.1. Let $0 < \alpha < \epsilon$ be two very small real numbers. Let $x_0 \in N$ and $A_0 = (x_0, y_0)$ the singular point of the projection map π_1 above x_0 . Let $x'_0 \in E \cap \mathbb{R}$ be a point such that $|x_0 - x'_0| < \alpha$. Then it is possible to enumerate the points of $K(x'_0) \cap \mathbb{R}$ and get that $K(x'_0) \cap \mathbb{R} = \{y_1 < \dots < y_n\}$. For sufficiently small ϵ there exists a unique pair of numbers k and l ($1 \leq k \leq l \leq n$) such that $K(x'_0) \cap (y_0 - \epsilon, y_0 + \epsilon) = \{y_k < \dots < y_l\}$. We call (k, l) the Lefschetz pair associated with the point x_0 and denote it by $\mathcal{L}(x_0)$.

Remark 4.2. Definition 4.1 does not hold when A_0 is of type d_2, f_2 and ν is even. In these cases we enumerate the real points of $K(x_0) = \{y_1 < \dots < y_n\}$, and take $k = l = j$ where y_j is the second coordinate of A_0 .

Remark 4.3. If A_0 is a point of type

- (1) $a_1, b, c, d_1, d_3, e_1, e_2, f_1, g_1, g_3$ we take $x'_0 > x_0$.
- (2) $a_2, d_2(\text{odd } \nu), f_2(\text{odd } \nu), g_2$ we take $x'_0 < x_0$.

Remark 4.4. If A_0 is a point of type

- (1) $a_1, a_2, b, d_1, d_2(\text{odd } \nu), e_1, e_2, g_3$ we have $l = k + 1$.
- (2) $f_1, f_2(\text{odd } \nu), g_1, g_2(\text{odd } \nu)$ we have $l = k + 2$.
- (3) c we have $l = k + n_l - 1$ where n_l is the number of intersecting (locally) non singular branches.
- (4) $d_3, d_2(\text{even } \nu), f_2(\text{even } \nu), g_2(\text{even } \nu)$ we have $l = k$.

Definition 4.5. Let $x_0 \in B$, and let $(x_0, y_0), (x_0, y'_0) \in S$ be the pair of points for which $0 < \Im(y_0) = \Im(y'_0)$. Take $x'_0 \in E \cap \mathbb{R}$ to be a point such that $|x'_0 - x_0| < \delta$ for a very small $0 < \delta$. Then it is possible to enumerate by the $\Im$ coordinate the points of $K(x'_0) \setminus \mathbb{R}$ and get $K(x'_0) \setminus \mathbb{R} = \{y_1 < \dots < y_d\}$. Since the curve S is defined over $\mathbb{R}$ we look at the set $A = \{\Im(y_1), \dots, \Im(y_d)\} \cap \mathbb{R}^+ = \{r_1, \dots, r_{\frac{d}{2}}\} = \{\Im(y_{\frac{d}{2}+1}), \dots, \Im(y_d)\}$. For sufficiently small $0 < \epsilon$ there exists a unique pair of numbers k and l ($1 \leq k \leq l \leq \frac{d}{2}$) such that $A \cap ((\Im(y_0) - \epsilon, \Im(y_0) + \epsilon)) = \{r_k, r_l\}$. We call the pair (k, l) the complex Lefschetz pair associated with the point x_0 and denote it by $\mathcal{L}(x_0)$.

Remark 4.6. Note that if $x_0 \in B$ and we have $\mathcal{L}(x_0) = (k, l)$ we always have $l = k + 1$. Moreover, if we take $A' = \{\Im(y_1), \dots, \Im(y_d)\} \cap \mathbb{R}^- = \{r_1, \dots, r_{\frac{d}{2}}\}$ in a decreasing order, we get an analogous definition to Definition 4.5, and the two complex Lefschetz pairs achieved from the two analogous definitions are the same.

Definition 4.7. Let $x_0 \in B$ with complex Lefschetz pair $\mathcal{L}(x_0) = (k, l)$, let $(x_0, y_0), (x_0, y'_0) \in S$ be the pairs of points for which $0 < \Im(y_0) = \Im(y'_0)$ and let A be as in Definition 4.5. Suppose that $\Re(y_0) < \Re(y'_0)$. Then, there exists a small enough $0 < \delta$ such that for every $x \in (x_0 - \delta, x_0 + \delta)$ if (x, y) is the lift of x to S which is connected to the point (x_0, y_0) and (x, y') is the lift of x to S which is connected to the point (x_0, y'_0) we have $\Re(y) < \Re(y')$, and δ is small enough to determine the complex Lefschetz pair.

Now, take $x'_0 \in E \cap \mathbb{R}$ such that $0 < x_0 - x'_0 < \delta$ and $x''_0 \in E \cap \mathbb{R}$ such that $0 < x''_0 - x_0 < \delta$. Denote by $\{z'_1, \dots, z'_{\frac{d}{2}}\} = \{z \in K(x'_0) \setminus \mathbb{R} \mid \Im(z) \in \mathbb{R}^+\}$ and by $\{z''_1, \dots, z''_{\frac{d}{2}}\} = \{z \in K(x''_0) \setminus \mathbb{R} \mid \Im(z) \in \mathbb{R}^+\}$.

We assign to x_0 an orientation $\mathcal{O}(x_0) \in \{1, -1, 0\}$ as follows:

- (1) If $\Re(z'_k) > \Re(z'_l)$ and $\Re(z''_k) < \Re(z''_l)$ then, $\mathcal{O}(x_0) = 1$.
- (2) If $\Re(z'_k) < \Re(z'_l)$ and $\Re(z''_k) > \Re(z''_l)$ then, $\mathcal{O}(x_0) = -1$.
- (3) Otherwise $\mathcal{O}(x_0) = 0$.

We are now going to use the model described in Section 3 in order to describe the algorithm.

Lemma 4.8. Let $\mathcal{R}$ be a connected component in $E \setminus (N \cup B)$, and suppose that we have $n - d$ real points in $K(x)$ for $x \in \mathcal{R}$. Then there exist a continuous family of diffeomorphisms $\{\beta_x \mid x \in \mathcal{R}\}$ such that:

- (1) $\beta_x : D \rightarrow \tilde{D}$
- (2) $\beta_x(K(x)) = K_d$
- (3) $\beta_x|_{\mathbb{R}} = \tilde{D} \cap \mathbb{R}$
- (4) For all $x, x' \in \mathcal{R}$ we have $\beta_x(y) = \beta_{x'}(y) \in \partial\tilde{D}$ for all $y \in \partial D$.

Proof. The proof follows immediately when noticing that for any two points x, x' there are exactly $n - d$ real points, and that since $\mathcal{R}$ does not contain any point in B the order in terms of $\Im$ coordinate of the non real points in $K(x)$ is fixed. Therefore, the points in $K(x)$ can be identified with the points in E , the real points of $K(x)$ with the points in K' , and the non real points of $K(x)$ with the points in K'' .

Now use a diffeomorphism which will transform the real points of $K(x)$ to K' maintaining the order described in Definition 4.1, the non real points of $K(x)$ to the non real points in $\tilde{D}$ ordered by their $\Im$ coordinate and fixing the image on ∂D . Since S is a curve, above $\mathcal{R}$ its elements are continuous so we can make sure that the family of diffeomorphisms is continuous as needed. $\square$

Remark 4.9. Although β defined in Lemma 4.8 is not unique, every two families of diffeomorphisms β and β' are homotopic. This fact will ensure us to have the correct result in the algorithm as we will describe later.

Definition 4.10. Let $s(t)$ be a curve in $E \setminus N$ parameterized by $t \in [0, 1]$. $s(t)$ has n lifts into S . Projecting these paths to D we get n paths in D starting at $K(s(0))$ and ending at $K(s(1))$. These paths induce a homotopy class of diffeomorphisms called the Lefschetz diffeomorphism induced by $s(t)$ and denoted by Ψ_s .

Proposition 4.11. Let $x_0 \in N$ and denote $A_0 = (x_0, y_0)$ the singular point of π_1 above x_0 . Let $s(t)$ be a parameterized semicircle of radius $0 < \alpha$ (very small) below the real line centered at x_0 , where $s(0) = -\alpha$ and $s(1) = \alpha$. Let Ψ_s be the Lefschetz diffeomorphism induced by s , and $\mathcal{L} = (k, l)$ be the Lefschetz pair associated with

x_0 . Then,

$$\beta_{x_0-\alpha}^{-1} \circ \Psi_s \circ \beta_{x_0+\alpha} = \begin{cases} \Delta_{a_1}^{\frac{1}{2}} \langle k, l \rangle & \text{if } A_0 \text{ is of type } a_1 \\ \Delta_{a_2}^{\frac{1}{2}} \langle k, l \rangle & \text{if } A_0 \text{ is of type } a_2 \\ \Delta_b^2 \langle k, l \rangle & \text{if } A_0 \text{ is of type } b \\ \Delta_c \langle k, l \rangle & \text{if } A_0 \text{ is of type } c \\ \Delta_{d_1}^{\frac{\nu}{2}} \langle k, l \rangle & \text{if } A_0 \text{ is of type } d_1 \\ \Delta_{d_2}^{\frac{2}{2}} \langle k \rangle & \text{if } A_0 \text{ is of type } d_2 \\ \Delta_{d_3}^{\frac{2}{3}} \langle k \rangle & \text{if } A_0 \text{ is of type } d_3 \\ \Delta_{e_1}^{\frac{1}{3}} \langle k, l \rangle & \text{if } A_0 \text{ is of type } e_1 \\ \Delta_{e_2}^{\frac{1}{3}} \langle k, l \rangle_{\langle a < b \rangle} & \text{if } A_0 \text{ is of type } e_2 \text{ and } a < b \\ \Delta_{e_2}^{\frac{1}{3}} \langle k, l \rangle_{\langle a > b \rangle} & \text{if } A_0 \text{ is of type } e_2 \text{ and } a > b \\ \Delta_{f_1}^{\frac{1}{2}} \langle k, l \rangle & \text{if } A_0 \text{ is of type } f_1 \\ \Delta_{f_2}^{\frac{1}{2}} \langle k, l \rangle & \text{if } A_0 \text{ is of type } f_2 \\ \Delta_{g_1}^{\frac{2}{2}} \langle k, l \rangle & \text{if } A_0 \text{ is of type } g_1 \\ \Delta_{g_2}^{\frac{\nu}{2}} \langle k, l \rangle & \text{if } A_0 \text{ is of type } g_2 \\ \Delta_{g_3}^{\frac{2}{3}} \langle k, l \rangle & \text{if } A_0 \text{ is of type } g_3 \end{cases}$$

Proof. In order to simplify notations we assume that $D = \tilde{D}$, the equations of S are all simple, and that the model suitable above the point α is E_d as described in Section 3.

The proof for points of types a_1, a_2, b, c can be found in Proposition 1.2 of [9].

A_0 is of type d_1 :

By Remark 4.4 we know that $k+1 = l$. Moreover, the component of S passing through A_0 has equation $(y - (k + \frac{1}{2}))^2 - x^\nu = 0$, and the other components of S have equations $y = c$ where $c \in K_d \setminus \{k, k+1\} = \{1, \dots, k-1, k+2, \dots, n-d, n+1 \pm i, \dots, n+1 \pm \frac{d}{2}i\}$. Now when ν is odd, we have that E_d is the model suitable above the point α , and that E_{d+2} is the suitable model above the point $-\alpha$. In addition, $K(-\alpha) = \{1, \dots, k-1, k+2, \dots, n-d, n+1 \pm i, \dots, n+1 \pm \frac{d}{2}i\} \cup \{k + \frac{1}{2} \pm i\sqrt{\alpha^\nu}\}$ and $K(\alpha) = \{1, \dots, k-1, k+2, \dots, n-d, n+1 \pm i, \dots, n+1 \pm \frac{d}{2}i\} \cup \{k + \frac{1}{2} \pm \sqrt{\alpha^\nu}\}$. Therefore, $\beta_{-\alpha}^{-1} : (D, K_{d+2}) \rightarrow (D, K(-\alpha))$ is induced by the motion: $n+1 \pm i \mapsto k + \frac{1}{2} \pm i\sqrt{\alpha^\nu}$, and so β_α is induced by the motion: $k + \frac{1}{2} \pm \sqrt{\alpha^\nu} \mapsto k + \frac{1}{2} \pm \frac{1}{2}$ and $n+1 + ij \mapsto n+1 + i(j-1)$ where $j = 2, \dots, \frac{d+2}{2}$.

Above s , S is the union of n paths. Projecting them to D we get the following motions:

$$\begin{array}{ll} y = c & \text{where } t : 0 \mapsto 1 \text{ and } c \in K_d \setminus \{k, k+1\} \\ y = k + \frac{1}{2} + \sqrt{\alpha^\nu(e^{i\pi+i\pi t})^\nu} & \text{where } t : 0 \mapsto 1 \\ y = k + \frac{1}{2} - \sqrt{\alpha^\nu(e^{i\pi+i\pi t})^\nu} & \text{where } t : 0 \mapsto 1 \end{array}$$

Therefore, $\Psi_s : (D, K(-\alpha)) \rightarrow (D, K(\alpha))$ is induced from the motion of the points $k + \frac{1}{2} \pm i\sqrt{\alpha^\nu}$ along a circle of radius $\sqrt{\alpha^\nu}$ centered at $k + \frac{1}{2}$ counterclockwise by angle of $\frac{\nu\pi}{2}$ to the points $k + \frac{1}{2} \pm \sqrt{\alpha^\nu}$.

When ν is even, the model suitable above the point $-\alpha$ is E_d and so, $K(-\alpha) = K(\alpha) = \{1, \dots, k-1, k+2, \dots, n-d, n+1 \pm i, \dots, n+1 \pm \frac{d}{2}i\} \cup \{k + \frac{1}{2} \pm \sqrt{\alpha^\nu}\}$. Therefore, $\beta_{-\alpha}^{-1} : (D, K_d) \rightarrow (D, K(-\alpha))$ is induced by the motion: $k \mapsto k + \frac{1}{2} - \sqrt{\alpha^\nu}$ and $k+1 \mapsto k + \frac{1}{2} + \sqrt{\alpha^\nu}$. Moreover, $\beta_\alpha = \beta_{-\alpha}$.

Above s , S is the union of n paths. Projecting them to D we get the following motions:

$$\begin{aligned} y &= c && \text{where } t : 0 \mapsto 1 \text{ and } c \in K_d \setminus \{k, k+1\} \\ y &= k + \frac{1}{2} + \sqrt{\alpha^\nu(e^{i\pi+i\pi t})^\nu} && \text{where } t : 0 \mapsto 1 \\ y &= k + \frac{1}{2} - \sqrt{\alpha^\nu(e^{i\pi+i\pi t})^\nu} && \text{where } t : 0 \mapsto 1 \end{aligned}$$

Therefore, $\Psi_s : (D, K(-\alpha)) \rightarrow (D, K(\alpha))$ is induced from the motion of the points $k + \frac{1}{2} \pm \sqrt{\alpha^\nu}$ along a circle of radius $\sqrt{\alpha^\nu}$ centered at $k + \frac{1}{2}$ counterclockwise by angle of $\frac{\nu\pi}{2}$ to the points $k + \frac{1}{2} \pm \sqrt{\alpha^\nu}$.

Hence, in any case of ν (odd or even), the composition of these motions is homotopic to the composition of the motions that induce $\Delta_{d_1}^{\frac{\nu}{2}} \langle k, l \rangle$. Therefore, the induced diffeomorphisms are homotopic, and we have $\beta_{x_0-\alpha}^{-1} \circ \Psi_s \circ \beta_{x_0+\alpha} = \Delta_{d_1}^{\frac{\nu}{2}} \langle k, l \rangle$.

A_0 is of type d_2 :

The proof is similar to the case where A_0 is of type d_1 .

A_0 is of type d_3 :

Since E_d is the model suitable above the point α , E_d is also the model suitable above the point $-\alpha$. The component of S passing through A_0 has equation $(y-k)^3 - x^2 = 0$, and other components of S have equations $y = c$ where $c \in K_d \setminus \{k, n+1 \pm i\} = \{1, \dots, k-1, k+1, \dots, n-d, n+1 \pm 2i, \dots, n+1 \pm \frac{d}{2}i\}$.

Now, $K(\alpha) = K(-\alpha) = \{1, \dots, k-1, k+1, \dots, n-d, n+1 \pm 2i, \dots, n+1 \pm i\frac{d}{2}\} \cup \{k + \sqrt[3]{\alpha^2}, k - \frac{1}{2}\sqrt[3]{\alpha^2} \pm i\frac{\sqrt{3}}{2}\sqrt[3]{\alpha^2}\}$.

$\beta_{-\alpha}^{-1} : (D, K_d) \rightarrow (D, K(-\alpha))$ is induced from the motion $n+1 \pm i \mapsto k - \frac{\sqrt[3]{\alpha^2}}{2} \pm i\frac{\sqrt{3}}{2}\sqrt[3]{\alpha^2}$, and $k \mapsto k + \sqrt[3]{\alpha^2}$. Moreover, $\beta_\alpha = \beta_{-\alpha}$.

Above s , S is the union of n paths. Projecting them to D we get the following motions:

$$\begin{aligned}
y = c & \quad \text{where } t : 0 \mapsto 1 \\
& \quad c \in K_d \setminus \{k, n+1 \pm i\} \\
y = k + (\alpha e^{i\pi+i\pi t})^{\frac{2}{3}} & \quad \text{where } t : 0 \mapsto 1 \\
y = k - \frac{1}{2}(\alpha e^{i\pi+i\pi t})^{\frac{2}{3}} + i\frac{\sqrt{3}}{2}(\alpha e^{i\pi+i\pi t})^{\frac{2}{3}} & \quad \text{where } t : 0 \mapsto 1 \\
y = k - \frac{1}{2}(\alpha e^{i\pi+i\pi t})^{\frac{2}{3}} - i\frac{\sqrt{3}}{2}(\alpha e^{i\pi+i\pi t})^{\frac{2}{3}} & \quad \text{where } t : 0 \mapsto 1
\end{aligned}$$

Therefore, $\Psi_s : (D, K(-\alpha)) \rightarrow (D, K(\alpha))$ is induced from the motion of the points $k + \sqrt[3]{\alpha^2}$ and $k - \frac{1}{2}\sqrt[3]{\alpha^2} \pm i\frac{\sqrt{3}}{2}\sqrt[3]{\alpha^2}$ along a circle of radius $\sqrt[3]{\alpha^2}$ centered at the point k counterclockwise by angle of $\frac{2\pi}{3}$ to themselves.

Hence, the composition of these three motions is homotopic to the composition of the motions that induce $\Delta_{d_3}^{\frac{2}{3}} \langle k \rangle$. Therefore, the induced diffeomorphisms are homotopic so we have $\beta_{x_0-\alpha}^{-1} \circ \Psi_s \circ \beta_{x_0+\alpha} = \Delta_{d_3}^{\frac{2}{3}} \langle k \rangle$.

A_0 is of type e_1 :

Since E_d is the model suitable above the point α it is also the model suitable above the point $-\alpha$. By Remark 4.4 we know that $k+1=l$. Moreover, the component of S passing through A_0 has equation $((y - (k + \frac{1}{2}))^4 - x^6 = 0$, and other components of S have equations $y = c$ where $c \in K_d \setminus \{k, k+1, n+1 \pm i\}$.

Now, $K(\alpha) = K(-\alpha) = \{1, \dots, k-1, k+2, \dots, n-d, n+1 \pm 2i, \dots, n+1 \pm i\frac{d}{2}\} \cup \{k + \frac{1}{2} \pm \sqrt{\alpha^3}, k + \frac{1}{2} \pm i\sqrt{\alpha^3}\}$.

$\beta_{-\alpha}^{-1} : (D, K_d) \rightarrow (D, K(-\alpha))$ is induced from the motion $k \mapsto k + \frac{1}{2} - \sqrt{\alpha^3}$, $k+1 \mapsto k + \frac{1}{2} + \sqrt{\alpha^3}$, $n+1 \pm i \mapsto k + \frac{1}{2} \pm i\sqrt{\alpha^3}$ in shortest lines. Moreover, $\beta_\alpha = \beta_{-\alpha}$.

Above s , S is the union of n paths. Projecting them to D we get the following motions:

$$\begin{aligned}
y = c & \quad \text{where } t : 0 \mapsto 1 \quad c \in K_d \setminus \{k, k+1, n+1 \pm i\} \\
y = k + \frac{1}{2} + \sqrt{(\alpha e^{i\pi+i\pi t})^3} & \quad \text{where } t : 0 \mapsto 1 \\
y = k + \frac{1}{2} - \sqrt{(\alpha e^{i\pi+i\pi t})^3} & \quad \text{where } t : 0 \mapsto 1 \\
y = k + \frac{1}{2} + i\sqrt{(\alpha e^{i\pi+i\pi t})^3} & \quad \text{where } t : 0 \mapsto 1 \\
y = k + \frac{1}{2} - i\sqrt{(\alpha e^{i\pi+i\pi t})^3} & \quad \text{where } t : 0 \mapsto 1
\end{aligned}$$

Therefore, $\Psi_s : (D, K(-\alpha)) \rightarrow (D, K(\alpha))$ is induced from the motion of the points $k + \frac{1}{2} \pm \sqrt{\alpha^3}$ and $k + \frac{1}{2} \pm i\sqrt{\alpha^3}$ along a circle of radius $\sqrt{\alpha^3}$ centered at $k + \frac{1}{2}$ counterclockwise by angle of $\frac{3\pi}{2}$ to themselves.

Hence, the composition of these three motions is homotopic to the composition of the motions that induce $\Delta_{e_1}^{\frac{3}{2}} \langle k, l \rangle$. Therefore, the induced diffeomorphisms are homotopic so we have $\beta_{x_0-\alpha}^{-1} \circ \Psi_s \circ \beta_{x_0+\alpha} = \Delta_{e_1}^{\frac{3}{2}} \langle k, l \rangle$.

A_0 is of type e_2 :

We deal with the two cases for points A_0 of type e_2 simultaneously. By Remark 4.4 we know that $k+1=l$, and we also know that the model suitable above the points α and $-\alpha$ is E_d . Moreover, the component of S passing through A_0 has equation $((y - (k + \frac{1}{2}))^3 + ax^2)((y - (k + \frac{1}{2}))^3 - bx^2) = 0$, and other components of S have equations $y = c$ where $c \in K_d \setminus \{k, k+1, n+1 \pm i, n+1 \pm 2i\}$.

Now, $K(\alpha) = K(-\alpha) = \{1, \dots, k-1, k+2, \dots, n-d, n+1 \pm 3i, \dots, n+1 \pm i\frac{d}{2}\} \cup \{k + \frac{1}{2} + \sqrt[3]{b\alpha^2}, k + \frac{1}{2} - \frac{\sqrt[3]{b\alpha^2}}{2} \pm i\frac{\sqrt{3}}{2}\sqrt[3]{b\alpha^2}, k + \frac{1}{2} - \sqrt[3]{a\alpha^2}, k + \frac{1}{2} + \frac{\sqrt[3]{a\alpha^2}}{2} \pm i\frac{\sqrt{3}}{2}\sqrt[3]{a\alpha^2}\}$

In order to know $\beta_{-\alpha}^{-1} : (D, K_d) \rightarrow (D, K(-\alpha))$ we need to separate between two cases:

$a > b$

$\beta_{-\alpha}^{-1} : (D, K_d) \rightarrow (D, K(-\alpha))$ is induced from the motion $k \mapsto k + \frac{1}{2} - \sqrt[3]{a\alpha^2}, k+1 \mapsto k + \frac{1}{2} + \sqrt[3]{b\alpha^2}, n+1 \pm i \mapsto k + \frac{1}{2} - \frac{1}{2}\sqrt[3]{b\alpha^2} \pm i\frac{\sqrt{3}}{2}\sqrt[3]{b\alpha^2}, n+1 \pm 2i \mapsto k + \frac{1}{2} + \frac{1}{2}\sqrt[3]{a\alpha^2} \pm i\frac{\sqrt{3}}{2}\sqrt[3]{a\alpha^2}$ in shortest lines.

$a < b$

$\beta_{-\alpha}^{-1} : (D, K_d) \rightarrow (D, K(-\alpha))$ is induced from the motion $k \mapsto k + \frac{1}{2} - \sqrt[3]{a\alpha^2}, k+1 \mapsto k + \frac{1}{2} + \sqrt[3]{b\alpha^2}, n+1 \pm i \mapsto k + \frac{1}{2} + \frac{1}{2}\sqrt[3]{a\alpha^2} \pm i\frac{\sqrt{3}}{2}\sqrt[3]{a\alpha^2}, n+1 \pm 2i \mapsto k + \frac{1}{2} - \frac{1}{2}\sqrt[3]{b\alpha^2} \pm i\frac{\sqrt{3}}{2}\sqrt[3]{b\alpha^2}$ in shortest lines.

Moreover, $\beta_\alpha = \beta_{-\alpha}$.

Above s , S is the union of n paths. Projecting them to D we get the following motions:

$y = c$	where $t : 0 \mapsto 1$ $c \in K_d \setminus \{k, k+1, n+1 \pm i, n+1 \pm 2i\}$
$k + \frac{1}{2} - \sqrt[3]{a\alpha^2}e^{2\pi it}$	where $t : 0 \mapsto 1$
$k + \frac{1}{2} + \sqrt[3]{b\alpha^2}e^{2\pi it}$	where $t : 0 \mapsto 1$
$k + \frac{1}{2} + \frac{1}{2}\sqrt[3]{a\alpha^2}e^{2\pi it} + i\frac{\sqrt{3}}{2}\sqrt[3]{a\alpha^2}e^{2\pi it}$	where $t : 0 \mapsto 1$
$k + \frac{1}{2} + \frac{1}{2}\sqrt[3]{a\alpha^2}e^{2\pi it} - i\frac{\sqrt{3}}{2}\sqrt[3]{a\alpha^2}e^{2\pi it}$	where $t : 0 \mapsto 1$
$k + \frac{1}{2} - \frac{1}{2}\sqrt[3]{b\alpha^2}e^{2\pi it} + i\frac{\sqrt{3}}{2}\sqrt[3]{b\alpha^2}e^{2\pi it}$	where $t : 0 \mapsto 1$
$k + \frac{1}{2} - \frac{1}{2}\sqrt[3]{b\alpha^2}e^{2\pi it} - i\frac{\sqrt{3}}{2}\sqrt[3]{b\alpha^2}e^{2\pi it}$	where $t : 0 \mapsto 1$

Therefore, $\Psi_s : (D, K(-\alpha)) \rightarrow (D, K(\alpha))$ is induced from the motion of the points $k + \frac{1}{2} - \sqrt[3]{a\alpha^2}, k + \frac{1}{2} + \sqrt[3]{b\alpha^2}, k + \frac{1}{2} + \frac{1}{2}\sqrt[3]{a\alpha^2} \pm i\frac{\sqrt{3}}{2}\sqrt[3]{a\alpha^2}, k + \frac{1}{2} - \frac{1}{2}\sqrt[3]{b\alpha^2} \pm i\frac{\sqrt{3}}{2}\sqrt[3]{b\alpha^2}$ centered at $k + \frac{1}{2}$ counterclockwise by angle of $\frac{2\pi}{3}$ to themselves.

The composition of these three motions is homotopic to the composition of the motions that induce $\Delta_{e_2}^{\frac{1}{3}} \langle k, l \rangle_{\langle a < b \rangle}$ or $\Delta_{e_2}^{\frac{1}{3}} \langle k, l \rangle_{\langle a > b \rangle}$ (depending whether $a < b$

or $b < a$). Therefore, the induced diffeomorphisms are homotopic and we have $\beta_{x_0-\alpha}^{-1} \circ \Psi_s \circ \beta_{x_0+\alpha} = \Delta_{e_2}^{\frac{1}{3}} \langle k, l \rangle$ (the $\langle a < b \rangle$ or $\langle b < a \rangle$) have been omitted to denote simultaneously both cases).

A_0 is of type f_1 :

By Remark 4.4 we know that $k+2 = l$. Moreover, the component of S passing through A_0 has equation $(y - (k+1))((y - (k+1))^2 - x^\nu) = 0$, and the other component of S have equations $y = c$ where $c \in K_d \setminus \{k, k+1, k+2\} = \{1, \dots, k-1, k+3, \dots, n-d, n+1 \pm i, \dots, n+1 \pm \frac{d}{2}i\}$.

Now, when ν is odd, we know that if the model suitable above the point α is E_d then E_{d+2} is the suitable model above the point $-\alpha$. In addition, $K(-\alpha) = \{1, \dots, k-1, k+3, \dots, n-d, n+1 \pm i, \dots, n+1 \pm \frac{d}{2}i\} \cup \{k+1, k+1 \pm i\sqrt{\alpha^\nu}\}$ and $K(\alpha) = \{1, \dots, k-1, k+3, \dots, n-d, n+1 \pm i, \dots, n+1 \pm \frac{d}{2}i\} \cup \{k+1, k+1 \pm \sqrt{\alpha^\nu}\}$. Therefore, $\beta_{-\alpha}^{-1} : (D, K_{d+2}) \rightarrow (D, K(-\alpha))$ is induced by the motion: $j \mapsto j+2$ where $j = k+1, \dots, n-d, k \mapsto k+1$ and $n+1 \pm i \mapsto k+1 \pm i\sqrt{\alpha^\nu}$. Moreover, β_α is induced by the motion: $k+1 \pm \sqrt{\alpha^\nu} \mapsto k+1 \pm 1$ and $n+1 + ij \mapsto n+1 + i(j-1)$ where $j = 2, \dots, \frac{d+2}{2}$.

Above s , S is the union of n paths. Projecting them to D we get the following motions:

$$\begin{array}{ll} y = c & \text{where } t : 0 \mapsto 1 \text{ and } c \in K_d \setminus \{k, k+1, k+2\} \\ y = k+1 & \text{where } t : 0 \mapsto 1 \\ y = k+1 + \sqrt{(\alpha e^{i\pi+i\pi t})^\nu} & \text{where } t : 0 \mapsto 1 \\ y = k+1 - \sqrt{(\alpha e^{i\pi+i\pi t})^\nu} & \text{where } t : 0 \mapsto 1 \end{array}$$

Therefore, $\Psi_s : (D, K(-\alpha)) \rightarrow (D, K(\alpha))$ is induced from the motion of the points $k+1 \pm i\sqrt{\alpha^\nu}$ along a circle of radius $\sqrt{\alpha^\nu}$ centered at $k+1$ counterclockwise by angle of $\frac{\nu\pi}{2}$ to the points $k+1 \pm \sqrt{\alpha^\nu}$.

When ν is even, we have that the model suitable above the points α and $-\alpha$ is E_d . $K(-\alpha) = K(\alpha) = \{1, \dots, k-1, k+3, \dots, n-d, n+1 \pm i, \dots, n+1 \pm \frac{d}{2}i\} \cup \{k+1, k+1 \pm \sqrt{\alpha^\nu}\}$.

Therefore, $\beta_{-\alpha}^{-1} : (D, K_d) \rightarrow (D, K(-\alpha))$ is induced by the motion: $k \mapsto k+1 - \sqrt{\alpha^\nu}$ and $k+2 \mapsto k+1 + \sqrt{\alpha^\nu}$. Moreover, $\beta_\alpha = \beta_{-\alpha}$.

Above s , S is the union of n paths. Projecting them to D we get the following motions:

$$\begin{array}{ll} y = c & \text{where } t : 0 \mapsto 1 \text{ and } c \in K_d \setminus \{k, k+1, k+2\} \\ y = k+1 & \text{where } t : 0 \mapsto 1 \\ y = k+1 + \sqrt{(\alpha e^{i\pi+i\pi t})^\nu} & \text{where } t : 0 \mapsto 1 \\ y = k+1 - \sqrt{(\alpha e^{i\pi+i\pi t})^\nu} & \text{where } t : 0 \mapsto 1 \end{array}$$

Therefore, $\Psi_s : (D, K(-\alpha)) \rightarrow (D, K(\alpha))$ is induced from the motion of the points $k+1 \pm \sqrt{\alpha^\nu}$ along a circle of radius $\sqrt{\alpha^\nu}$ centered at $k+1$ counterclockwise by angle of $\frac{\nu\pi}{2}$ to the points $k+1, k+\frac{1}{2} \pm \sqrt{\alpha^\nu}$.

Hence, in any case of ν (odd or even), the composition of these motions is homotopic to the composition of the motions that induce $\Delta_{f_1}^{\frac{\nu}{2}} \langle k, l \rangle$. Therefore, the induced diffeomorphisms are homotopic. We have $\beta_{x_0-\alpha}^{-1} \circ \Psi_s \circ \beta_{x_0+\alpha} = \Delta_{f_1}^{\frac{\nu}{2}} \langle k, l \rangle$.

A_0 is of type f_2 :

The proof is similar to the case where A_0 is of type f_1 .

A_0 is of type g_1 :

By Remark 4.4 we know that $k+2=l$. Moreover, the component of S passing through A_0 has equation $(y-(k+1)-ax)((y-(k+1))^2-x^\nu)=0$, where $0 < a$ and the other components of S have equations $y=c$ where $c \in K_d \setminus \{k, k+1, k+2\} = \{1, \dots, k-1, k+3, \dots, n-d, n+1 \pm i, \dots, n+1 \pm \frac{d}{2}i\}$.

Now when ν is odd, we know that since the model suitable above the point α is E_d , the model suitable above the point $-\alpha$ is E_{d+2} . In addition, $K(-\alpha) = \{1, \dots, k-1, k+3, \dots, n-d, n+1 \pm i, \dots, n+1 \pm \frac{d}{2}i\} \cup \{k+1-a\alpha, k+1 \pm i\sqrt{\alpha^\nu}\}$ and $K(\alpha) = \{1, \dots, k-1, k+3, \dots, n-d, n+1 \pm i, \dots, n+1 \pm \frac{d}{2}i\} \cup \{k+1+a\alpha, k+1 \pm \sqrt{\alpha^\nu}\}$.

Therefore, $\beta_{-\alpha}^{-1} : (D, K_{d+2}) \rightarrow (D, K(-\alpha))$ is induced by the motion: $j \mapsto j+2$ where $j = k+1, \dots, n-d, k \mapsto k+1-a\alpha$ and $n+1 \pm i \mapsto k+1 \pm i\sqrt{\alpha^\nu}$. Moreover, β_α is induced by the motion: $k+1 \pm \sqrt{\alpha^\nu} \mapsto k+\frac{1}{2} \pm \frac{1}{2}, k+1+a\alpha \mapsto k+2$ and $n+1 \pm ij \mapsto n+1 \pm i(j-1)$ where $j = 2, \dots, \frac{d}{2}$.

Above s , S is the union of n paths. Projecting them to D we get the following motions:

$$\begin{array}{ll} y = c & \text{where } t : 0 \mapsto 1 \text{ and } c \in K_d \setminus \{k, k+1, k+2\} \\ y = k+1 + a\alpha e^{i\pi+i\pi t} & \text{where } t : 0 \mapsto 1 \\ y = k+1 + \sqrt{(\alpha e^{i\pi+i\pi t})^\nu} & \text{where } t : 0 \mapsto 1 \\ y = k+1 - \sqrt{(\alpha e^{i\pi+i\pi t})^\nu} & \text{where } t : 0 \mapsto 1 \end{array}$$

Therefore, $\Psi_s : (D, K(-\alpha)) \rightarrow (D, K(\alpha))$ is induced from the motion of the points $k+1 \pm i\sqrt{\alpha^\nu}$ along a circle of radius $\sqrt{\alpha^\nu}$ centered at $k+1$ counterclockwise by angle of $\frac{\nu\pi}{2}$ to the points $k+1 \pm \sqrt{\alpha^\nu}$, and the motion of the point $k+1-a\alpha$ to the point $k+1+a\alpha$ along a circle of radius $a\alpha$ counterclockwise by angle of π .

When ν is even, we have that E_d is the model suitable for both points α and $-\alpha$. $K(-\alpha) = K(\alpha) = \{1, \dots, k-1, k+3, \dots, n-d, n+1 \pm i, \dots, n+1 \pm \frac{d}{2}i\} \cup \{k+1-a\alpha, k+1 \pm \sqrt{\alpha^\nu}\}$.

Therefore, $\beta_{-\alpha}^{-1} : (D, K_d) \rightarrow (D, K(-\alpha))$ is induced by the motion: $k \mapsto k+1-a\alpha$, $k+1 \mapsto k+1-\sqrt{\alpha^\nu}$ and $k+2 \mapsto k+1+\sqrt{\alpha^\nu}$. Moreover, $\beta_\alpha = \beta_{-\alpha}$.

Above s , S is the union of n paths. Projecting them to D we get the following motions:

$$\begin{aligned} y &= c && \text{where } t : 0 \mapsto 1 \text{ and } c \in K_d \setminus \{k, k+1, k+2\} \\ y &= k+1 + a\alpha e^{i\pi+i\pi t} && \text{where } t : 0 \mapsto 1 \\ y &= k+1 + \sqrt{(\alpha e^{i\pi+i\pi t})^\nu} && \text{where } t : 0 \mapsto 1 \\ y &= k+1 - \sqrt{(\alpha e^{i\pi+i\pi t})^\nu} && \text{where } t : 0 \mapsto 1 \end{aligned}$$

Therefore, $\Psi_s : (D, K(-\alpha)) \rightarrow (D, K(\alpha))$ is induced from the motion of the points $k+1 \pm \sqrt{\alpha}^\nu$ along a circle of radius $\sqrt{\alpha}^\nu$ centered at $k+1$ counterclockwise by angle of $\frac{\nu\pi}{2}$ to themselves, and the motion of the point $k+1 - a\alpha$ to the point $k+1 + a\alpha$ along a circle of radius $a\alpha$ centered at $k+1$ counterclockwise by angle of π .

Hence, in any case of ν (odd or even), the composition of these motions is homotopic to the composition of the motions that induce $\Delta_{g_1}^{\frac{\nu}{2}} \langle k, l \rangle$. Therefore, the induced diffeomorphisms are homotopic. We have $\beta_{x_0-\alpha}^{-1} \circ \Psi_s \circ \beta_{x_0+\alpha} = \Delta_{g_1}^{\frac{\nu}{2}} \langle k, l \rangle$.

A_0 is of type g_2 :

The proof is similar to the case where A_0 is of type g_1 .

A_0 is of type g_3 :

By Remark 4.4 we know that $k+1 = l$. We also know that the model suitable above both points α and $-\alpha$ is E_d . Moreover, the component of S passing through A_0 has equation $(y-k)((y-k)^3 - x^2) = 0$, and other components of S have equations $y = c$ where $c \in K_d \setminus \{k, k+1, k+2, k+3\} = \{1, \dots, k-1, k+4, \dots, n-d, n+1 \pm i, \dots, n+1 \pm \frac{d}{2}i\}$.

Now, $K(\alpha) = K(-\alpha) = \{1, \dots, k-1, k+4, \dots, n-d, n+1 \pm i, \dots, n+1 \pm i\frac{d}{2}\} \cup \{k, k + \sqrt[3]{\alpha^2}, k - \frac{1}{2}\sqrt[3]{\alpha^2} \pm i\frac{\sqrt[3]{3}}{2}\sqrt[3]{\alpha^2}\}$.

$\beta_{-\alpha}^{-1} : (D, K_d) \rightarrow (D, K(-\alpha))$ is induced from the motion $n+1 \pm i \mapsto k - \frac{1}{2}\sqrt[3]{\alpha^2} \pm i\frac{\sqrt[3]{3}}{2}\sqrt[3]{\alpha^2}$, $j \mapsto j+2$ where $j = k+2, \dots, n-d$ and $k+1 \mapsto k + \sqrt[3]{\alpha^2}$ in shortest lines. Moreover, $\beta_\alpha = \beta_{-\alpha}$.

Above s , S is the union of n paths. Projecting them to D we get the following motions:

$$\begin{aligned} y &= c \text{ where } t : 0 \mapsto 1 && c \in K_d \setminus \{k, k+1, k+2, k+3\} \\ y &= k + \sqrt[3]{(\alpha e^{i\pi+i\pi t})^2} && \text{where } t : 0 \mapsto 1 \\ y &= (k - \frac{1}{2}\sqrt[3]{(\alpha e^{i\pi+i\pi t})^2} + i\frac{\sqrt[3]{3}}{2}\sqrt[3]{(\alpha e^{i\pi+i\pi t})^2}) && \text{where } t : 0 \mapsto 1 \\ y &= (k - \frac{1}{2}\sqrt[3]{(\alpha e^{i\pi+i\pi t})^2} - i\frac{\sqrt[3]{3}}{2}\sqrt[3]{(\alpha e^{i\pi+i\pi t})^2}) && \text{where } t : 0 \mapsto 1 \\ y &= k && \text{where } t : 0 \mapsto 1 \end{aligned}$$

Therefore, $\Psi_s : (D, K(-\alpha)) \rightarrow (D, K(\alpha))$ is induced from the motion of the points $k, k + \sqrt[3]{\alpha^2}$ and $(k - \frac{1}{2}\sqrt[3]{\alpha^2} \pm i\frac{\sqrt{3}}{2}\sqrt[3]{\alpha^2})$ along a circle of radius $\sqrt[3]{\alpha^2}$ centered at k counterclockwise by angle of $\frac{2\pi}{3}$ to themselves.

The composition of these three motions is homotopic to the composition of the motions that induce $\Delta_{g_3}^{\frac{1}{3}} \langle k, l \rangle$. Therefore, the induced diffeomorphisms are homotopic so we have $\beta_{x_0-\alpha}^{-1} \circ \Psi_s \circ \beta_{x_0+\alpha} = \Delta_{g_3}^{\frac{1}{3}} \langle k, l \rangle$.

□

Proposition 4.12. *Let $x_0 \in B$. Let u be a parameterized (from $x_0 - \alpha$ to $x_0 + \alpha$) the straight line segment $[x_0 - \alpha, x_0 + \alpha]$ (α very small), and let Ψ_u be the Lefschetz diffeomorphism induced by u , and $\mathcal{L} = (k, l)$ be the complex Lefschetz pair associated with x_0 . Then,*

$$\Delta_{com}^{\mathcal{O}(x_0)} \langle k, l \rangle = \beta_{x_0-\alpha}^{-1} \Psi_u \beta_{x_0+\alpha}$$

Proof. First it follows from Remark 4.6 that the diffeomorphism must be symmetrical along the real axis; Therefore, we concentrate on the half of the complex plane which consists of the points with positive imaginary coordinate, keeping in mind that what ever happens there also happens in the other half-plane in a mirror like image. Recall from Definition 4.5 that if $(x_0, y_0), (x_0, y'_0) \in S$ is one of the pairs of points for which $0 < \Im(y_0) = \Im(y'_0)$ and A as in Definition 4.5 we take small enough $0 < \alpha$ such that the complex Lefschetz pair is well defined and that for every $x \in (x_0 - \alpha, x_0 + \alpha)$ if (x, y) is the lift of x to S which is connected to the point (x_0, y_0) and (x, y') is the lift of x to S which is connected to the point (x_0, y'_0) we have $\Re(y) < \Re(y')$ when $\Re(y_0) < \Re(y'_0)$ or $\Re(y) > \Re(y')$ when $\Re(y_0) > \Re(y'_0)$ (It can't be that $\Re(y_0) = \Re(y'_0)$ or else the point x_0 will be in N instead of B). Without lose of generality we assume that $\Re(y_0) < \Re(y'_0)$.

Now as in Definition 4.7, let $x'_0 \in E \cap \mathbb{R}$ such that $0 < x_0 - x'_0 < \alpha$ and $x''_0 \in E \cap \mathbb{R}$ such that $0 < x''_0 - x_0 < \alpha$. Denote by $\{z'_1, \dots, z'_{\frac{d}{2}}\} = \{z \in K(x') \setminus \mathbb{R} \mid \Im(z) \in \mathbb{R}^+\}$ and by $\{z''_1, \dots, z''_{\frac{d}{2}}\} = \{z \in K(x'') \setminus \mathbb{R} \mid \Im(z) \in \mathbb{R}^+\}$.

$\beta_{x_0-\alpha}^{-1} : (D, K_d) \rightarrow (D, K(\alpha))$ is induced by the motion of the point $n+1+ki \mapsto z'_k, n+1+li \mapsto z'_l$ and the points $n+1-ki, n+1-li$ goes to the complex conjugates of z'_k and z'_l along straight lines (we assume that all other components of S have equations $y = c$ where $c \in K_d \setminus \{n+1 \pm ik, n+1 \pm il\}$).

$\beta_{x_0+\alpha} : (D, K(\alpha)) \rightarrow (D, K_d)$ is induced by the motion of the points z''_k, z''_l and their complex conjugates to the points $n+1 \pm ki, n+1 \pm li$ along straight lines.

We look at the following cases:

$\Re(z'_k) > \Re(z'_l)$ and $\Re(z''_k) < \Re(z''_l)$

In this case the point (x'_0, z'_k) is connected to the point (x_0, y'_0) and the point (x'_0, z'_l) is connected to the point (x_0, y_0) . Because $k < l$ we know by Lemma 4.8 that $\Im(z'_k) < \Im(z'_l)$. Now, since $\Re(z''_k) < \Re(z''_l)$ the point (x''_0, z''_k) is connected to the point (x_0, y_0) and the point (x''_0, z''_l) is connected to the point (x_0, y'_0) . However, $\Im(z''_k) < \Im(z''_l)$ hence we have a motion in which the point on the left goes down and the point on the right goes up crossing each other $\Im$ coordinate once (see Figure 9 (left)). Therefore, the composition $\beta_{x_0-\alpha}^{-1} \Psi_u \beta_{x_0+\alpha}$ is homotopic to the diffeomorphism defining $\Delta_{com}^1 \langle k, l \rangle$ (see Figure 9 (right)).




FIGURE 9. Positive motion in the fiber (left), $\beta_{x_0-\alpha}^{-1} \Psi_u \beta_{x_0+\alpha}$ (right)

$\Re(z'_k) < \Re(z'_l)$ and $\Re(z''_k) > \Re(z''_l)$ In this case the point (x'_0, z'_k) is connected to the point (x_0, y_0) and the point (x'_0, z'_l) is connected to the point (x_0, y'_0) . Because $k < l$ we know by Lemma 4.8 that $\Im(z'_k) < \Im(z'_l)$. Now, since $\Re(z''_k) > \Re(z''_l)$ the point (x''_0, z''_k) is connected to the point (x_0, y'_0) and the point (x''_0, z''_l) is connected to the point (x_0, y_0) . However, $\Im(z''_k) < \Im(z''_l)$ hence we have a motion in which the point on the left goes up and the point on the right goes down crossing each other $\Im$ coordinate once (see Figure 10 (left)). Therefore, the composition $\beta_{x_0-\alpha}^{-1} \Psi_u \beta_{x_0+\alpha}$ is homotopic to the diffeomorphism defining $\Delta_{com}^{-1} \langle k, l \rangle$ (see Figure 10 (right)).




FIGURE 10. Negative motion in the fiber (left), $\beta_{x_0-\alpha}^{-1} \Psi_u \beta_{x_0+\alpha}$ (right)

Otherwise In these cases we have either $\Re(z'_k) < \Re(z'_l)$ and $\Re(z''_k) < \Re(z''_l)$ or $\Re(z'_k) > \Re(z'_l)$ and $\Re(z''_k) > \Re(z''_l)$. Hence the point on the left and on the right move until their $\Im$ coordinates coincide and then retract back. In this case there

is no crossing therefore the diffeomorphism induced by the motion which induce $\beta_{x_0-\alpha}^{-1}\Psi_u\beta_{x_0+\alpha}$ is trivial as can be seen in Figure 11 (one of two possible cases).




FIGURE 11. Trivial motion in the fiber (left), $\beta_{x_0-\alpha}^{-1}\Psi_u\beta_{x_0+\alpha}$ (right)

□

We need to be able to describe the braids of the local braid monodromy at each point of the set N in a way that will make it easy to conjugate them. This can be achieved by using a product of half-twists that define the resulted braid. This product is the parallel of the skeleton definition given in [8] and [9].

Definition 4.13. Let $\tilde{K} \subset \hat{K} \subset \tilde{D}$. A skeleton in $(\tilde{D}, \hat{K}, \tilde{K})$ is a consecutive sequence of paths each starting and ending in one of the points of $\tilde{K}$. Each path starts where it's previous ends. A skeleton is denoted by ξ .

Definition 4.14. We call a product of half-twists that describes the local braid monodromy at the point $x_0 \in N$ the generalized skeleton of x_0 , and denote it by ξ_{x_0} .

Recall that half-twists are defined by paths, so in order to describe the proper generalized skeleton for each type of singularity, it is enough to describe the paths defining its half-twists, and give their product form.

Definition 4.15. Let $x_0 \in N$, and $A_0 = (x_0, y_0)$ be the singular point of π_1 above x_0 . Let $\mathcal{L}_{x_0} = (k, l)$ be the Lefschetz pair associated with the point x_0 . We define the generalized skeletons:

- (1) If A_0 is of singularity type a_1 , the generalized skeleton assigned to it is h_1 , where h_1 is described in Figure 12 (a).
- (2) If A_0 is of singularity type a_2 , the generalized skeleton assigned to it is h_1 , where h_1 is described in Figure 12 (b).
- (3) If A_0 is of singularity type b , the generalized skeleton assigned to it is h_1^2 , where h_1 is described in Figure 12 (a).
- (4) If A_0 is of singularity type c , the generalized skeleton assigned to it is $(h_1 \cdots h_{l-k})(h_1 \cdots h_{l-k-1}) \cdots (h_1 h_2)(h_1)$, where h_i are described in Figure 12 (c).

- (5) If A_0 is of singularity type d_1 , the generalized skeleton assigned to it is h_1^ν , where h_1 is described in Figure 12 (a).
- (6) If A_0 is of singularity type d_2 , the generalized skeleton assigned to it is h_1^ν , where h_1 is described in Figure 12 (b).
- (7) If A_0 is of singularity type d_3 , the generalized skeleton assigned to it is $(h_1 h_2)^2$, where h_i are described in Figure 12 (d).
- (8) If A_0 is of singularity type e_1 , the generalized skeleton assigned to it is $(h_2 h_1 h_3)^6$, where h_1, h_2, h_3 are described in Figure 12 (e).
- (9) If A_0 is of singularity type e_2 , the generalized skeleton assigned to it is $h_2 h_4 h_1 h_3 h_5$, where h_i are described in Figure 12 (f) when $b < a$, and in Figure 12 (g) when $a < b$.
- (10) If A_0 is of singularity type f_1 , the generalized skeleton assigned to it is $(h_1 h_2 h_1)^\nu$, where h_1, h_2 are described in Figure 12 (c).
- (11) If A_0 is of singularity type f_2 , the generalized skeleton assigned to it is $(h_2 h_1 h_2)^\nu$, where h_1, h_2 are described in Figure 12 (h).
- (12) If A_0 is of singularity type g_1 , the generalized skeleton assigned to it is $h_2 h_1 h_2^\nu h_1 h_2$, where h_1, h_2 are described in Figure 12 (c).
- (13) If A_0 is of singularity type g_2 , the generalized skeleton assigned to it is $h_2 h_1 h_2^\nu h_1 h_2$, where h_1 is described in Figure 12 (h).
- (14) If A_0 is of singularity type g_3 , the generalized skeleton assigned to it is $(h_1 h_2 h_3 h_2)^\nu$, where h_1, h_2, h_3 are described in Figure 12 (i).




FIGURE 12. Half-twists used in products defining generalized skeletons

Theorem 4.16. *Let $x_0 \in N$, and $A_0 = (x_0, y_0)$ be the singular point of π_1 above x_0 . Let $\mathcal{L}_{x_0} = (k, l)$ be the Lefschetz pair associated with the point x_0 . Then, the generalized skeletons defined in Definition 4.15, describe the local braid monodromy at the point x_0 .*

Proof. The proof follows immediately from Proposition 2.1 to Proposition 2.9 and the proofs in [9]. $\square$

4.2. Description of a generating set for $\pi_1(E \setminus N)$.

We need to compute the braid monodromy homomorphism. Recall that φ is a homomorphism from $\pi_1(E \setminus N)$, which is the fundamental group of a punctured disk, to the braid group B_n . Therefore, in order to describe the mapping, we choose a generating set for $\pi_1(E \setminus N)$, and compute the image of φ on these elements. The image of φ on this generating set not only describe completely the braid monodromy homomorphism but also serve as an important step in the computation of the fundamental group of the complement of the curve.

We fix as a base point for the fundamental group of the origin of φ the point x_0 which is the larger of the two points from $\partial E \cap \mathbb{R}$. Moreover we enumerate the points of N such that $\{x_1 > \dots > x_p\}$. Then, we take an ordered system of paths below the real axis. Each of the paths circles exactly one point in N , as can be seen in Figure 13. We denote the elements of this generating set by $\Gamma_1, \dots, \Gamma_p$.




FIGURE 13. The generating set for $\pi_1(E \setminus N)$

Now we turn to describe these paths.

To each point $x_j \in N$ we attach a real value $0 < \alpha_j$ which is not only sufficiently small in order to determine the Lefschetz pair $\mathcal{L}(x_j) = (k_j, l_j)$ for the point x_j but also small enough such that $(x_j - \alpha_j, x_j + \alpha_j) \cap (N \cup B) = \{x_j\}$. Denote by t_0 the

real segment between $x_1 + \alpha_1$ and x_0 , and for any $1 \leq j < p$, denote $t_j(t)$ $t \in [0, 1]$ the real segment $[x_{j+1} + \alpha_{j+1}, x_j - \alpha_j]$.

Denote also by $s_j(t)$ the parametrization of a semi circle below the real axis centered at x_j with radius α_j , and $r_j(t)$ the full circle oriented counterclockwise centered at x_j with radius α_j for $t \in [0, 1]$, such that $s_j(0) = x_j - \alpha_j$ and $s_j(1) = x_j + \alpha_j$, and $r_j(0) = r_j(1) = x_j + \alpha_j$. We also denote the type of singularity of the singular point of π_1 above x_j by p_j (i.e., $p_j \in \{a_1, a_2, b, c, d_1, d_2, d_3, e_1, e_2, f_1, f_2, g_1, g_2, g_3\}$).

The closed disk E with the notations above can be seen in Figure 14.




FIGURE 14. $\pi_1(E \setminus N)$ with notations

Γ_j is now homotopic to the following path:

$$\Gamma_j = t_0^{-1} \left(\prod_{k=1}^{j-1} (s_k^{-1} t_k^{-1}) \right) r_j \left(\prod_{k=j-1}^1 (t_k s_k) \right) t_0.$$

We take a closer look at the segment t_k . Denote the ordered set of points $t_k \cap B = \{b_{k,1} > \dots > b_{k,n_k}\}$ the points of the set B in the segment t_k . To each of these points, using definition 4.5, we attached a positive small real number $0 < \delta_{k,l}$ where $l = 1, \dots, n_k$. The set of points $\{x_k - \alpha_k, b_{k,1} + \delta_{k,1}, b_{k,1} - \delta_{k,1}, \dots, b_{k,n_k} + \delta_{k,n_k}, b_{k,n_k} - \delta_{k,n_k}, x_{k+1} + \alpha_{k+1}\}$ induce a partition on t_k . Here too, we take $\delta_{k,l}$ (and α) small enough such that $(b_{k,l} - \delta_{k,l}, b_{k,l} + \delta_{k,l}) \cap (N \cup B) = \{b_{k,l}\}$, and $(b_{k,l} - \delta_{k,l}, b_{k,l} + \delta_{k,l}) \cap (x_j - \alpha_j, x_j + \alpha_j) = \emptyset$ For all k, l, j .

Denote:

- (1) $t_{k,0} = [b_{k,1} + \delta_{k,1}, x_k - \alpha_k] \forall k \neq 0$.
- (2) $t_{k,l} = [b_{k,l+1} + \delta_{k,l+1}, b_{k,l} - \delta_{k,l}] \forall l = 1, \dots, n_k - 1$.
- (3) $t_{k,n_k} = [x_{k+1} + \alpha_{k+1}, b_{k,n_k} - \delta_{k,n_k}]$.
- (4) $u_{k,l} = [b_{k,l} - \delta_{k,l}, b_{k,l} + \delta_{k,l}] \forall l = 1, \dots, n_k$.
- (5) $t_{0,0} = [b_{0,1} + \delta_{0,1}, x_0]$.

This partition implies that:

$$t_k = \left(\prod_{l=n_k}^1 t_{k,l} u_{k,l} \right) t_{k,0}.$$

Hence,

$$\Gamma_j = \left(t_{0,0}^{-1} \left(\prod_{l=1}^{n_0} u_{0,l}^{-1} t_{0,l}^{-1} \right) \right) \prod_{k=1}^{j-1} (s_k^{-1} t_{k,0}^{-1} \left(\prod_{l=1}^{n_k} u_{k,l}^{-1} t_{k,l}^{-1} \right)) r_j \prod_{k=j-1}^1 \left(\left(\prod_{l=n_k}^1 t_{k,l} u_{k,l} \right) t_{k,0} s_k \right) \left(\left(\prod_{l=n_0}^1 t_{0,l} u_{0,l} \right) t_{0,0} \right).$$

4.3. The computation of $\varphi(\Gamma_j)$. Using the definitions above it follows that if

$$\begin{aligned} \mathcal{Q} &= \left(\prod_{k=j-1}^1 \Psi_{t_k} \Psi_{s_k} \right) \Psi_{t_0}. \\ &= \prod_{k=j-1}^1 \left(\left(\prod_{l=n_k}^1 \Psi_{t_{k,l}} \Psi_{u_{k,l}} \right) \Psi_{t_{k,0}} \Psi_{s_k} \right) \left(\prod_{l=n_0}^1 \Psi_{t_{0,l}} \Psi_{u_{0,l}} \right) \Psi_{t_{0,0}}. \end{aligned}$$

Then, the Lefschetz diffeomorphism of Γ_j is the product:

$$\Psi_{\Gamma_j} = \mathcal{Q}^{-1} \Psi_{r_j} \mathcal{Q}$$

which implies that

$$\varphi(\Gamma_j) = \mathcal{Q}^{-1} \Delta_{p_j} \mathcal{Q}.$$

Where Δ_{p_j} is the local braid monodromy defined in Section 3, which complies with the topological type of the singular point x_j denoted p_j , and with the Lefschetz pair associated with it $\mathcal{L}(x_j)$.

In the model defined above this is realized as follows:

$$\begin{aligned} \Psi_{\Gamma_j} &= \beta_{x_0}^{-1} \mathcal{Q}^{-1} \Delta_{p_j} \mathcal{Q} \beta_{x_0} \\ &= (\beta_{x_0}^{-1} \mathcal{Q}^{-1} \beta_{x_j+\alpha_j}) (\beta_{x_j+\alpha_j}^{-1} \Delta_{p_j} \beta_{x_j+\alpha_j}) (\beta_{x_j+\alpha_j}^{-1} \mathcal{Q} \beta_{x_0}). \end{aligned}$$

We turn now to $\beta_{x_j+\alpha_j}^{-1} \mathcal{Q} \beta_{x_0}$.

$$\begin{aligned} \beta_{x_j+\alpha_j}^{-1} \mathcal{Q} \beta_{x_0} &= \beta_{x_j+\alpha_j}^{-1} \left(\prod_{k=j-1}^1 \Psi_{t_k} \Psi_{s_k} \right) \Psi_{t_0} \beta_{x_0} \\ &= \left(\prod_{k=j-1}^1 (\beta_{x_{k+1}+\alpha_{k+1}}^{-1} \Psi_{t_k} \beta_{x_k-\alpha_k}) (\beta_{x_k-\alpha_k}^{-1} \Psi_{s_k} \beta_{x_k+\alpha_k}) \right) (\beta_{x_1+\alpha_1}^{-1} \Psi_{t_0} \beta_{x_0}) \end{aligned}$$

By Proposition 4.11 we know that $\beta_{x_k-\alpha_k}^{-1} \Psi_{s_k} \beta_{x_k+\alpha_k} = \Delta_{p_k} \langle k, l \rangle$ where p_k is the singularity type as defined above, and the power is omitted. Therefore, we need to study $\beta_{x_{k+1}+\alpha_{k+1}}^{-1} \Psi_{t_k} \beta_{x_k+\alpha_k}$. However, t_k is actually a product so if one identifies the point $b_{k,n_k+1} + \delta_{k,n_k+1}$ as the point $x_{k+1} + \alpha_{k+1}$ and takes $\alpha_0 = 0$ we have:

$$\begin{aligned}
\beta_{x_{k+1}+\alpha_{k+1}}^{-1} \Psi_{t_k} \beta_{x_k-\alpha_k} &= \beta_{x_{k+1}+\alpha_{k+1}}^{-1} \left(\prod_{l=n_k}^1 \Psi_{t_{k,l}} \Psi_{u_{k,l}} \right) \Psi_{t_{k,0}} \beta_{x_k-\alpha_k} \\
&= \left(\prod_{l=n_k}^1 (\beta_{b_{k,l+1}+\delta_{k,l+1}}^{-1} \Psi_{t_{k,l}} \beta_{b_{k,l}-\delta_{k,l}}) (\beta_{b_{k,l}-\delta_{k,l}}^{-1} \Psi_{u_{k,l}} \beta_{b_{k,l}+\delta_{k,l}}) \right) \\
&\quad (\beta_{b_{k,1}+\delta_{k,1}} \Psi_{t_{k,0}} \beta_{x_k-\alpha_k}).
\end{aligned}$$

However, by Lemma 4.8 we have that the following four diffeomorphisms all induce the trivial braid:

- (1) $\beta_{b_{0,1}+\delta_{0,1}}^{-1} \Psi_{t_{0,0}} \beta_{x_0}$
- (2) $\beta_{b_{k,l+1}+\delta_{k,l+1}}^{-1} \Psi_{t_{k,l}} \beta_{b_{k,l}-\delta_{k,l}}$ for $k = 0, \dots, p-1$ and $l = 1, \dots, n_j - 1$.
- (3) $\beta_{b_{k,1}+\delta_{k,1}}^{-1} \Psi_{t_{k,0}} \beta_{x_k-\alpha_k}$ for $k = 1, \dots, p-2$.
- (4) $\beta_{x_{k+1}+\alpha_{k+1}}^{-1} \Psi_{t_{k,n_k}} \beta_{b_{k,n_k}-\delta_{k,n_k}}$ for $k = 1, \dots, p-1$.

Therefore, by Propositions 4.11 and 4.12 as a braid we get the following diffeomorphism of the model:

$$\begin{aligned}
&\beta_{x_j+\alpha_j}^{-1} \mathcal{Q} \beta_{x_0} \\
&= \prod_{k=j-1}^1 \left(\left(\prod_{l=n_k}^1 (\beta_{b_{k,l}-\delta_{k,l}}^{-1} \Psi_{u_{k,l}} \beta_{b_{k,l}+\delta_{k,l}}) \right) (\beta_{x_k-\alpha_k}^{-1} \Psi_{s_k} \beta_{x_k+\alpha_k}) \right) \left(\prod_{l=n_0}^1 (\beta_{b_{0,l}-\delta_{0,l}}^{-1} \Psi_{u_{0,l}} \beta_{b_{0,l}+\delta_{0,l}}) \right) \\
&= \prod_{k=j-1}^1 \left(\left(\prod_{l=n_k}^1 \Delta_{com}^{\mathcal{O}(b_{k,l})} \langle k_{k,l}, l_{k,l} \rangle \right) \Delta_{p_k}^{e_k} \langle k_{x_k}, l_{x_k} \rangle \right) \left(\prod_{l=n_0}^1 \Delta_{com}^{\mathcal{O}(b_{0,l})} \langle k_{0,l}, l_{0,l} \rangle \right)
\end{aligned}$$

where $\Delta_{p_k}^{e_k} \langle k_{x_k}, l_{x_k} \rangle$ is the diffeomorphism in Proposition 4.11.

Recall that by Definition 4.15 each Δ_{p_j} is a generalized skeleton which is a product of half-twists. Therefore, it is possible to write $\Delta_{x_j} = \prod_m (H_m)$ where H_m are the half-twists from Definition 4.15.

Moreover, it was proved in [8] that on a half-twist conjugation is done by activating on the path defining the half-twist the diffeomorphism conjugating it. Therefore, we can calculate the braid monodromy $\varphi(\Gamma_j)$ using the following:

Theorem 4.17. *Let $S, N, x_j, p_j, \mathcal{L}(x_j) = \langle k_{x_j}, l_{x_j} \rangle, \mathcal{L}(b_{k,l}) = \langle k_{k,l}, l_{k,l} \rangle, e_k, \alpha_j$ and $\delta_{k,l}$ be as above. Then,*

$$\begin{aligned}
\varphi(\Gamma_j) &= \prod_m \left(\prod_{k=j-1}^1 \left(\left(\prod_{l=n_k}^1 \Delta_{com}^{\mathcal{O}(b_{k,l})} \langle k_{k,l}, l_{k,l} \rangle \right) \Delta_{p_k}^{e_k} \langle k_{x_k}, l_{x_k} \rangle \right) \left(\prod_{l=n_0}^1 \Delta_{com}^{\mathcal{O}(b_{0,l})} \langle k_{0,l}, l_{0,l} \rangle \right) \right) (H_m) \\
&\quad (\text{as a braid}).
\end{aligned}$$

□

Remark 4.18. *It is always possible to choose the point x_0 in such a way that $B \cap [x_1, x_0] = \emptyset$. Hence, we may assume that in t_0 there are no points that belong to the set B .*

Remark 4.19. *The exact location of the points in the set $N \cup B$ is not important for the algorithm. What is important, however, is the order of the points with respect to each other.*

4.4. The algorithm. Now when the theory was established we can formulate the algorithm for computing the braid monodromy of a curve S . It's main idea is to find all the critical points which effect the braid monodromy results, the singular points of the curve and the points where complex switching occur. Then, using the computed information take each element of the generalized skeleton and compute for the result of the action induced by the diffeomorphisms conjugations as formulated in Theorem 4.17. The product of the results as a product of braids is the braid monodromy for this generator of the fundamental group. The ordered set $\varphi(\Gamma_j)$ ($j = 1, \dots, n$) define, as a product, the *braid monodromy factorization* which describes the braid monodromy homomorphism.

Algorithm 4.20. *Algorithm for computing the braid monodromy of a curve S*

- (1) *Find in S all the singular points with respect to the generic projection π_1 .*
- (2) *Compute for each point found in step 1 its projection to the first coordinate. The set of all projections is the set N .*
- (3) *Compute for each point $x_j \in N$ its Lefschetz pair $\mathcal{L}(x_j) = (k_{x_j}, l_{x_j})$.*
- (4) *Find for each $x_j \in N$ the diffeomorphism $\Delta_{p_j}^{e_j}$ induced by x_j on the model by Proposition 4.11.*
- (5) *Find in S all the complex switching points with respect to π_1 .*
- (6) *Compute for each point found in step 4 its projection to the first coordinate. The set of all projections is the set B .*
- (7) *Compute for each point $b_{k,l} \in B$ its complex Lefschetz pair $\mathcal{L}(b_{k,l}) = (k_{k,l}, l_{k,l})$, and find the orientation $\mathcal{O}(b_{k,l})$ induced by the point using Definition 4.7.*
- (8) *For each point x_j find using Definition 4.15 the generalized skeleton $\prod_m H_m$ defining its local braid monodromy.*
- (9) *For each point x_j compute the product which is the braid monodromy result on Γ_j using the generalized skeleton from Step 7 and Theorem 4.17:*

$$\varphi(\Gamma_j) = \prod_m \left(\prod_{k=j-1}^1 \left(\left(\prod_{l=n_k}^1 \Delta_{com}^{\mathcal{O}(b_{k,l})} \langle k_{k,l}, l_{k,l} \rangle \right) \Delta_{p_k}^{e_k} \langle k_{x_k}, l_{x_k} \rangle \right) \left(\prod_{l=n_0}^1 \Delta_{com}^{\mathcal{O}(b_{0,l})} \langle k_{0,l}, l_{0,l} \rangle \right) \right) (H_m)$$

In the next Section we give examples for the computation of braid monodromy using the algorithm.

5. TWO EXAMPLES FOR THE ALGORITHM USE

5.1. The computation of the braid monodromy of a cusp. In order to demonstrate the algorithm we compute the braid monodromy for the image of the curve S defined by the equation $y^2 - x^3 = 0$ under a linear transformation. In order to properly modify the curve using a linear transformation we take it's homogenization to $\mathbb{CP}^2$ which is defined by the homogenous polynomial: $y^2w - x^3$.

Then, we use the matrix $\begin{pmatrix} 5 & 3 & 1 \\ 3 & 2 & 1 \\ 2 & 1 & 1 \end{pmatrix}$ defining a linear transformation, to look at

the image of the curve under this transformation, which is the curve defined by $(3x + 2y + w)^2(2x + y + w) - (5x + 3y + w)^3 = 0$. In order to compute it's braid monodromy we choose generically a line at infinity and look at the curve's affine part which is $(3x + 2y + 1)^2(2x + y + 1) - (5x + 3y + 1)^3 = 0$.

Following steps 1 and 2 of Algorithm 4.20, we find the points of the set N . Figure 15 shows the real part of the curve with it's singular points identified. This also serve us in Step 2 so we know the order of the projections to the first coordinate of the singular points of S .




FIGURE 15. The curve defined by $(3x + 2y + 1)^2(2x + y + 1) - (5x + 3y + 1)^3 = 0$.

By Step 3, we compute the Lefschetz pair for each one of the singular points. $\mathcal{L}(x_1) = (2, 3)$, $\mathcal{L}(x_2) = (1, 2)$, $\mathcal{L}(x_3) = (1, 2)$, $\mathcal{L}(x_4) = (2, 3)$.

In Step 4, we find the diffeomorphisms induced by each point depending on it's Lefschetz pair and singularity type: $\Delta_{a_2}^{\frac{1}{2}} \langle 2, 3 \rangle$, $\Delta_{d_1}^{\frac{3}{2}} \langle 1, 2 \rangle$, $\Delta_{a_2}^{\frac{1}{2}} \langle 1, 2 \rangle$, $\Delta_{a_1}^{\frac{1}{2}} \langle 2, 3 \rangle$ for the points x_1, x_2, x_3, x_4 respectively.

Note that since our curve is of degree 3, we could have at most 2 conjugate complex points and therefore, $B = \emptyset$. Hence, Step 5,6 and 7 of the algorithm are trivial.

In Step 8, we find the generalized skeleton for each singular point and so we may summarize what we have until now in Table 1.

Point	Type	$\mathcal{L}$	$\prod_m H_m$	Δ_{x_j}
x_1	a_2	$\mathcal{L}(x_1) = \langle 2, 3 \rangle$	$\begin{array}{c} \cdot \\ \downarrow \end{array}$	$\Delta_{a1}^{1/2} \langle 2, 3 \rangle$
x_2	d_1	$\mathcal{L}(x_2) = \langle 1, 2 \rangle$	$\begin{array}{c} \leftarrow 3 \rightarrow \cdot \end{array}$	$\Delta_{d1}^{3/2} \langle 1, 2 \rangle$
x_3	a_2	$\mathcal{L}(x_3) = \langle 1, 2 \rangle$	$\begin{array}{c} \curvearrowright \cdot \end{array}$	$\Delta_{a1}^{1/2} \langle 1, 2 \rangle$
x_4	a_1	$\mathcal{L}(x_4) = \langle 2, 3 \rangle$	$\begin{array}{c} \cdot \longrightarrow \end{array}$	$\Delta_{a1}^{1/2} \langle 2, 3 \rangle$

TABLE 1. Results of steps 1–8 of Algorithm 4.20 on the curve $(3x + 2y + 1)^2(2x + y + 1) - (5x + 3y + 1)^3 = 0$

Finally, we compute the braid monodromy using Step 9:

Note, that all generalized skeletons consists of only one half-twist (besides the generalized skeleton of the point x_2 which is a half-twist taken to the power 3). Now, for each j we need to conjugate the generalized skeleton of x_j by all $\Delta_{p_k}^{e_k} \langle \mathcal{L}(x_k) \rangle$, where $k = j - 1, \dots, 1$. This is best summarized Table 3 (See Appendix B). Each part of table is used for the computation of the braid monodromy of a different Γ_j . First line in the is the generalized skeleton half-twists determined by the singular point which it's product represents the local braid monodromy at the point x_j . The left side of each line (besides the first) is the diffeomorphism which acts on the generalized skeleton elements, where the right side of each line is the result of this action. Finally, taking the product of the generalized skeleton elements at the last line is the result of the braid monodromy of Γ_j .

If the braid group of the model of the fiber above x_0 is generated by the two half-twists σ_1 and σ_2 as shown in Figure 16, we conclude that:

$$\begin{aligned} \varphi(\Gamma_1) &= \sigma_2 \\ \varphi(\Gamma_2) &= \sigma_1^3 \end{aligned}$$



FIGURE 16. σ_1 (left) and σ_2 (right), the generators of B_3 in the model above x_0 .

$$\begin{aligned}\varphi(\Gamma_3) &= \sigma_1 \\ \varphi(\Gamma_4) &= \sigma_1^{-2} \sigma_2 \sigma_1^2\end{aligned}$$

5.2. Example for the use of the algorithm when $B \neq \emptyset$. For the second example consider the curve S defined by the polynomial $(x^2 - (y + \frac{3}{4})^2 - 1)((x - \frac{1}{2})^2 - y^2 - 1)$. We follow the steps of Algorithm 4.20 in order to compute it's braid monodromy.

In Step 1, we compute all the singular points with respect to the projection π_1 , and in Step 2, we project these points to get the set N . In our case the set N consists of 6 points as can be seen in Figure 17.




FIGURE 17. The real part of the curve defined by $(x^2 - (y + \frac{3}{4})^2 - 1)((x - \frac{1}{2})^2 - y^2 - 1)$.

Next step in the algorithm (3) is to compute the Lefschetz pair associated with each point found in Steps 1 and 2. We get $\mathcal{L}(x_1) = (3, 4)$, $\mathcal{L}(x_2) = (2, 3)$, $\mathcal{L}(x_3) = (1, 2)$, $\mathcal{L}(x_4) = (1, 2)$, $\mathcal{L}(x_5) = (2, 3)$, $\mathcal{L}(x_6) = (1, 2)$.

In Step 4, we find the diffeomorphisms induced by each singular point: $\Delta_c \langle 2, 3 \rangle$, $\Delta_{a_1}^{\frac{1}{2}} \langle 2, 3 \rangle$, $\Delta_{a_1}^{\frac{1}{2}} \langle 1, 2 \rangle$, $\Delta_{a_2}^{\frac{1}{2}} \langle 1, 2 \rangle$, $\Delta_{a_2}^{\frac{1}{2}} \langle 2, 3 \rangle$, $\Delta_c \langle 1, 2 \rangle$ for the points $x_1, \dots, x_6$ respectively.

By Step 5, we find all the complex switching points with respect to π_1 . This can be done for example by solving the equations of the curve to y , isolating the $\Im$ coordinate, and finding real x for which we have equalities. It is also evident from Figure 5.2 which shows only the $\Im$ coordinates of the equation above the real axis, that we have one point in B which is positioned between x_3 and x_4 . Hence, we have completed Step 5 and 6 and we have the set $B = \{b_{3,1} = b\}$. Moreover, since there are at most 4 complex points at any fiber above the real axis, we know that the complex Lefschetz pair associated with the point b is $\mathcal{L}(b) = \langle 1, 2 \rangle$. As for the orientation induced by the point b we follow Definition 4.7. To the right of b the points $n + 1 \pm i$ in the model belong to the imaginary points with smaller real coordinate, and the points $n + 1 \pm 2i$ belong to the imaginary points with larger real coordinate. To the left of b the points $n + 1 \pm i$ in the model belong to the imaginary points with larger real coordinate, and the points $n + 1 \pm 2i$ in the model belong to the imaginary points with the smaller real coordinate. Hence, the orientation induced by b is $\mathcal{O}(b) = 1$, and so we have completed Step 7 of the algorithm.




FIGURE 18. The $\Im$ coordinates of the curve defined by $(x^2 - (y + \frac{3}{4})^2 - 1)((x - \frac{1}{2})^2 - y^2 - 1)$ above the real axis.

In Step 8, we find the generalized skeletons induced by each of the singular points found in Step 1. Results of the steps of the algorithm up to now are summarized in Table 2.

Finally, we compute the braid monodromy using Step 9:


For each x_j we need to conjugate the elements constructing the generalized skeleton

Point	type	$\mathcal{L}$	$\prod_m H_m$	Δ_{x_j}
x_1	c	$\mathcal{L}(x_1) = \langle 3, 4 \rangle$	$\cdot \cdot \xrightarrow{2}$	$\Delta_c \langle 3, 4 \rangle$
x_2	a_1	$\mathcal{L}(x_2) = \langle 2, 3 \rangle$	$\cdot \xrightarrow{1} \cdot$	$\Delta_{a_1}^{\frac{1}{2}} \langle 2, 3 \rangle$
x_3	a_1	$\mathcal{L}(x_3) = \langle 1, 2 \rangle$	$\xrightarrow{1} \cdot$	$\Delta_{a_1}^{\frac{1}{2}} \langle 1, 2 \rangle$
b	com	$\mathcal{L}(b) = \langle 1, 2 \rangle$	No generalized skeleton	$\Delta_{com}^1 \langle 1, 2 \rangle$
x_4	a_2	$\mathcal{L}(x_4) = \langle 1, 2 \rangle$	$\cdot$ $\downarrow$ $\cdot$	$\Delta_{a_2}^{\frac{1}{2}} \langle 1, 2 \rangle$
x_5	a_2	$\mathcal{L}(x_5) = \langle 2, 3 \rangle$	$\cdot \cdot \xrightarrow{2}$	$\Delta_{a_2}^{\frac{1}{2}} \langle 2, 3 \rangle$
x_6	c	$\mathcal{L}(x_6) = \langle 1, 2 \rangle$	$\xrightarrow{2} \cdot \cdot$	$\Delta_c \langle 1, 2 \rangle$

TABLE 2. Results of steps 1–8 of Algorithm 4.20 on the curve $(x^2 - (y + \frac{3}{4})^2 - 1)((x - \frac{1}{2})^2 - y^2 - 1)$.

of x_j by all $\Delta_{p_k}^{e_k} \langle \mathcal{L}(x_k) \rangle$, where $k = j-1, \dots, 1$, and p_k is the singularity type of the point x_k or the diffeomorphism induced by the point b . This is best summarized Table 4 (See Appendix B). In order to demonstrate the use of the algorithm we bring here only full computations for the points x_5 and x_6 , for the rest of the points we give only the final result.

If the braid group of the model of the fiber above x_0 is generated by the three half-twists σ_1, σ_2 and σ_3 as shown in Figure 19, we conclude that:
 $\varphi(\Gamma_1) = \sigma_3^2$





FIGURE 19. The generators of B_4 in the model above x_0





$$\begin{aligned}
 \varphi(\Gamma_2) &= \sigma_3^{-1} \sigma_2 \sigma_3 \\
 \varphi(\Gamma_3) &= \sigma_2^{-1} \sigma_1 \sigma_2 \\
 \varphi(\Gamma_4) &= \sigma_2^{-1} \sigma_3 \sigma_2 \\
 \varphi(\Gamma_5) &= \sigma_3^2 \sigma_2^{-1} \sigma_1 \sigma_2 \sigma_3^{-2} \\
 \varphi(\Gamma_6) &= \sigma_3^2
 \end{aligned}$$






REFERENCES






- [1] M. Amram, D. Garber and M. Teicher., *Local braid monodromies and local fundamental groups of tangented conic-line arrangements*, submitted (math.GT/0401351).
- [2] M. Amram, D. Garber and M. Teicher., *Fundamental groups of tangented conic-line arrangements with singularities up to order 4*, submitted (math.GT/0305418).
- [3] E. Artin, *Theory of braids*, Ann. Math. **48** (1947), 101-126.
- [4] J. Birman, *Braids, links and mapping class groups*, Ann. Math Studies 82, Princeton University Press, 1975.
- [5] O. Chisini, *Sulla identità birazionale delle funzioni algebriche di due variabili dotate di una medesima curva di diramazione*, Rend. Ist. Lombardo **77**, 339–356 (1944).
- [6] D. Cohen and A. Suciu., *The braid monodromy of plane algebraic curves and hyperplane arrangements*, Commentarii Math. Helvetici **72** (1997), 285-315.
- [7] P. Dehornoy, *Braids and Self Distributivity*, Progress in Mathematics, **192**; xvi + 624 pages. Birkhauser (2000)
- [8] B. Moishezon and M. Teicher, *Braid group techniques in complex geometry I, Line arrangements in $\mathbb{CP}^2$* , Contemporary Math. **78** (1988), 425-555.
- [9] B. Moishezon and M. Teicher, *Braid Group Techniques in Complex Geometry II: From arrangements of lines to cuspidal curves*, LNM **1479** (1989), 131-179.
- [10] V. S. Kulikov, *On Chisini's conjecture*, Izv. Ross. Akad. Nauk Ser. Mat. **63**:6, 83-116 (1999) [Russian]; *English transl.*, Izvestiya Math. **63**, 1139-1170 (1999).
- [11] V. S. Kulikov and M. Teicher, *Braid monodromy factorizations and diffeomorphism types*, Izv. Ross. Akad. Nauk Ser. Mat. **64**(2), 89-120 (2000) [Russian]; *English transl.*, Izvestiya Math. **64**(2), 311-341 (2000).

6. APPENDIX A


 FIGURE 20. $\Delta_{a_1}^{\frac{1}{2}} \langle k, l \rangle$

 FIGURE 21. $\Delta_{a_2}^{\frac{1}{2}} \langle k, l \rangle$

 FIGURE 22. $\Delta_b^2 \langle k, l \rangle$

 FIGURE 23. $\Delta_c \langle k, l \rangle$

FIGURE 24. $\Delta_{d_1}^{\frac{\nu}{2}} \langle k, l \rangle$ with ν oddFIGURE 25. $\Delta_{d_1}^{\frac{\nu}{2}} \langle k, l \rangle$ with ν evenFIGURE 26. $\Delta_{d_2}^{\frac{\nu}{2}}$ with ν oddFIGURE 27. $\Delta_{d_2}^{\frac{\nu}{2}}$ with ν even


 FIGURE 28. $\Delta_{d_3}^{\frac{2}{3}} \langle k, l \rangle$

 FIGURE 29. $\Delta_{e_1}^{\frac{3}{2}} \langle k, l \rangle$

 FIGURE 30. $\Delta_{e_2}^{\frac{2}{3}} \langle k, l \rangle_{\langle a < b \rangle}$

 FIGURE 31. $\Delta_{e_2}^{\frac{2}{3}} \langle k, l \rangle_{\langle a > b \rangle}$

 FIGURE 32. $\Delta_{f_1}^{\frac{\nu}{2}} \langle k, l \rangle$ with ν odd

FIGURE 33. $\Delta_{f_1}^{\frac{\nu}{2}} \langle k, l \rangle$ with ν evenFIGURE 34. $\Delta_{f_2}^{\frac{\nu}{2}} \langle k, l \rangle$ with ν oddFIGURE 35. $\Delta_{f_2}^{\frac{\nu}{2}} \langle k, l \rangle$ with ν evenFIGURE 36. $\Delta_{g_1}^{\frac{\nu}{2}} \langle k, l \rangle$ with ν oddFIGURE 37. $\Delta_{g_1}^{\frac{\nu}{2}} \langle k, l \rangle$ with ν even





 FIGURE 38. $\Delta_{g_2}^{\frac{\nu}{2}} \langle k, l \rangle$ with ν odd

 FIGURE 39. $\Delta_{g_2}^{\frac{\nu}{2}} \langle k, l \rangle$ with ν even

 FIGURE 40. $\Delta_{g_3}^{\frac{2}{3}} \langle k, l \rangle$

 FIGURE 41. $\Delta_{com}^1 \langle k, l \rangle$

7. APPENDIX B

$\varphi(\Gamma_1)$		$\varphi(\Gamma_2)$	
Conj. diffeo.	Generalized skeleton	Conj. diffeo.	Generalized skeleton
		$\Delta_{a1}^{1/2} < 2, 3 >$	
$\varphi(\Gamma_3)$		$\varphi(\Gamma_4)$	
Conj. diffeo.	Generalized skeleton	Conj. diffeo.	Generalized skeleton
$\Delta_{d1}^{3/2} < 1, 2 >$		$\Delta_{a1}^{1/2} < 1, 2 >$	
$\Delta_{a1}^{1/2} < 2, 3 >$		$\Delta_{d1}^{3/2} < 1, 2 >$	
		$\Delta_{a1}^{1/2} < 2, 3 >$	

TABLE 3. Braid monodromy for the cusp defined by the polynomial $(3x + 2y + 1)^2(2x + y + 1) - (5x + 3y + 1)^3$

$\varphi(\Gamma_1)$		$\varphi(\Gamma_2)$	
Conj. diffeo.	Generalized skeleton	Conj. diffeo.	Generalized skeleton
Final result		Final result	
$\varphi(\Gamma_3)$		$\varphi(\Gamma_4)$	
Conj. diffeo.	Generalized skeleton	Conj. diffeo.	Generalized skeleton
Final result		Final result	
$\varphi(\Gamma_5)$		$\varphi(\Gamma_6)$	
Conj. diffeo.	Generalized skeleton	Conj. diffeo.	Generalized skeleton
$\Delta_{a_2}^{\frac{1}{2}} \langle 1, 2 \rangle$		$\Delta_{a_2}^{\frac{1}{2}} \langle 2, 3 \rangle$	
$\Delta_{com} \langle 1, 2 \rangle$		$\Delta_{a_2}^{\frac{1}{2}} \langle 1, 2 \rangle$	
$\Delta_{a_1}^{\frac{1}{2}} \langle 1, 2 \rangle$		$\Delta_{com} \langle 1, 2 \rangle$	
$\Delta_{a_1}^{\frac{1}{2}} \langle 2, 3 \rangle$		$\Delta_{a_1}^{\frac{1}{2}} \langle 1, 2 \rangle$	
$\Delta_c \langle 3, 4 \rangle$		$\Delta_{a_1}^{\frac{1}{2}} \langle 2, 3 \rangle$	
		$\Delta_c \langle 3, 4 \rangle$	

TABLE 4. Braid monodromy for the curve defined by the polynomial $(x^2 - (y + \frac{3}{4})^2 - 1)((x - \frac{1}{2})^2 - y^2 - 1)$

DEPARTMENT OF MATHEMATICS AND STATISTICS, BAR-ILAN UNIVERSITY, RAMAT-GAN
52900, ISRAEL

E-mail address: `kaplansh@macs.biu.ac.il`

DEPARTMENT OF MATHEMATICS AND STATISTICS, BAR-ILAN UNIVERSITY, RAMAT-GAN
52900, ISRAEL

E-mail address: `liberme1@macs.biu.ac.il`

DEPARTMENT OF MATHEMATICS AND STATISTICS, BAR-ILAN UNIVERSITY, RAMAT-GAN
52900, ISRAEL

E-mail address: `teicher@macs.biu.ac.il`